

ANTARES Constrains a Blazar Origin of Two IceCube PeV Neutrino Events

S. Adrián-Martínez¹, A. Albert², M. André³, G. Anton⁵, M. Ardid¹,
 J.-J. Aubert⁶, B. Baret⁷, J. Barrios⁸, S. Basa⁹, V. Bertin⁶, S. Biagi²³,
 C. Bogazzi¹², R. Bormuth^{12,13}, M. Bou-Cabo¹, M.C. Bouwhuis¹²,
 R. Bruijn^{12,14}, J. Brunner⁶, J. Busto⁶, A. Capone^{15,16}, L. Caramete¹⁷,
 J. Carr⁶, T. Chiarusi¹⁰, M. Circella¹⁸, R. Coniglione²³, L. Core⁶,
 H. Costantini⁶, P. Coyle⁶, A. Creusot⁷, G. De Rosa^{19,20}, I. Dekeyser²¹,
 A. Deschamps²², G. De Bonis^{15,16}, C. Distefano²³, C. Donzaud^{7,24},
 D. Dornic⁶, Q. Dorosti²⁵, D. Drouhin², A. Dumas²⁶, T. Eberl⁵,
 A. Enzenhöfer⁵, S. Escoffier⁶, K. Fehn⁵, I. Felis¹, P. Fermani^{15,16},
 F. Folger⁵, L.A. Fusco^{10,11}, S. Galatà⁷, P. Gay²⁶, S. Geißelsöder⁵,
 K. Geyer⁵, V. Giordano²⁷, A. Gleixner⁵, J.P. Gómez-González⁸,
 K. Graf⁵, G. Guillard²⁶, H. van Haren²⁸, A.J. Heijboer¹², Y. Hello²²,
 J.J. Hernández-Rey⁸, B. Herold⁵, A. Herrero¹, J. Höfl⁵,
 J. Hofestädt⁵, C. Hugon⁴, C.W James^{* 5}, M. de Jong^{12,13},
 O. Kalekin⁵, U. Katz⁵, D. Kießling⁵, P. Kooijman^{12,14,29},
 A. Kouchner⁷, V. Kulikovskiy²³, R. Lahmann⁵, E. Lambard⁶,
 G. Lambard⁸, D. Lefèvre²¹, E. Leonora^{27,32}, H. Loehner²⁵,
 S. Loucatos³³, S. Mangano⁸, M. Marcelin⁹, A. Margiotta^{10,11},
 J.A. Martínez-Mora¹, S. Martini²¹, A. Mathieu⁶, T. Michael¹²,
 P. Migliozzi¹⁹, M. Neff⁵, E. Nezri⁹, D. Palioselitis¹², G.E. Pāvāļš¹⁷,
 C. Pellegrino^{10,11}, C. Perrina^{15,16}, P. Piattelli²³, V. Popa¹⁷,
 T. Pradier³⁴, C. Racca², G. Riccobene²³, R. Richter⁵, K. Roensch⁵,
 A. Rostovtsev³⁵, M. Saldaña¹, D. F. E. Samtleben^{12,13},
 A. Sánchez-Losa⁸, M. Sanguineti^{4,36}, P. Sapienza²³, J. Schmid⁵,
 J. Schnabel⁵, S. Schulte¹², F. Schüssler³³, T. Seitz⁵, C. Sieger⁵,
 A. Spies⁵, M. Spurio^{10,11}, J.J.M. Steijger¹², Th. Stolarczyk³³,
 M. Taiuti^{4,36}, C. Tamburini²¹, Y. Tayalati³⁷, A. Trovato²³,
 B. Vallage³³, C. Vallée⁶, V. Van Elewyck⁷, E. Visser¹², D. Vivolo^{19,20},
 S. Wagner⁵, E. de Wolf^{12,14}, K. Yatkin⁶, H. Yepes⁸, J.D. Zornoza⁸,
 J. Zúñiga⁸,
 The TANAMI Collaboration: F. Krauß^{30,38}, M. Kadler^{†38},
 K. Mannheim³⁸, R. Schulz^{30,38}, J. Trüstedt^{30,38}, J. Wilms³⁰,

R. Ojha^{39,40,41}, E. Ros^{42,43,44}, W. Baumgartner³⁹, T. Beuchert^{30,38},
 J. Blanchard⁴⁵, C. Bürkel^{30,38}, B. Carpenter⁴¹, P.G. Edwards⁴⁶,
 D. Eisenacher Glawion³⁸, D. Elsässer³⁸, U. Fritsch⁵, N. Gehrels³⁹,
 C. Gräfe^{30,38}, C. Großberger⁴⁷, H. Hase⁴⁸, S. Horiuchi⁴⁹, A. Kappes³⁸,
 A. Kreikenbohm^{30,38}, I. Kreykenbohm³⁰, M. Langejahn^{30,38},
 K. Leiter^{30,38}, E. Litzinger^{30,38}, J.E.J. Lovell⁵⁰, C. Müller^{30,38},
 C. Phillips⁴⁶, C. Plötz⁴⁸, J. Quick⁵¹, T. Steinbring^{30,38}, J. Stevens⁴⁶,
 D. J. Thompson³⁹, and A.K. Tzioumis⁴⁶

- ¹Institut d'Investigació per a la Gestió Integrada de les Zones Costaneres (IGIC) - Universitat Politècnica de València. C/ Paranimf 1, 46730 Gandia, Spain
- ²GRPHE - Université de Haute Alsace - Institut Universitaire de Technologie de Colmar, 34 rue du Grillenbreit BP 50568 - 68008 Colmar, France
- ³Technical University of Catalonia, Laboratory of Applied Bioacoustics, Rambla Exposició, 08800 Vilanova i la Geltrú, Barcelona, Spain
- ⁴INFN - Sezione di Genova, Via Dodecaneso 33, 16146 Genova, Italy
- ⁵Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen Centre for Astroparticle Physics, Erwin-Rommel-Str. 1, 91058 Erlangen, Germany
- ⁶CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France
- ⁷APC, Université Paris Diderot, CNRS/IN2P3, CEA/IRFU, Observatoire de Paris, Sorbonne Paris Cité, 75205 Paris, France
- ⁸IFIC - Instituto de Física Corpuscular, Edificios Investigación de Paterna, CSIC - Universitat de València, Apdo. de Correos 22085, 46071 Valencia, Spain
- ⁹LAM - Laboratoire d'Astrophysique de Marseille, Pôle de l'Étoile Site de Château-Gombert, rue Frédéric Joliot-Curie 38, 13388 Marseille Cedex 13, France
- ¹⁰INFN - Sezione di Bologna, Viale Berti-Pichat 6/2, 40127 Bologna, Italy
- ¹¹Dipartimento di Fisica e Astronomia dell'Università, Viale Berti-Pichat 6/2, 40127 Bologna, Italy
- ¹²Nikhef, Science Park, Amsterdam, The Netherlands
- ¹³Huygens-Kamerlingh Onnes Laboratorium, Universiteit Leiden, The Netherlands
- ¹⁴Universiteit van Amsterdam, Instituut voor Hoge-Energie Fysica, Science Park 105, 1098 XG Amsterdam, The Netherlands
- ¹⁵INFN -Sezione di Roma, P.le Aldo Moro 2, 00185 Roma, Italy
- ¹⁶Dipartimento di Fisica dell'Università La Sapienza, P.le Aldo Moro 2, 00185 Roma, Italy
- ¹⁷Institute for Space Science, RO-077125 Bucharest, Măgurele, Romania
- ¹⁸INFN - Sezione di Bari, Via E. Orabona 4, 70126 Bari, Italy
- ¹⁹INFN -Sezione di Napoli, Via Cintia 80126 Napoli, Italy
- ²⁰Dipartimento di Fisica dell'Università Federico II di Napoli, Via Cintia 80126, Napoli, Italy
- ²¹Mediterranean Institute of Oceanography (MIO), Aix-Marseille University, 13288, Marseille, Cedex 9, France; Université du Sud Toulon-Var, 83957, La Garde Cedex, France
 CNRS-INSU/IRD UM 110
- ²²Géoazur, Université Nice Sophia-Antipolis, CNRS/INSU, IRD, Observatoire de la Côte d'Azur, Sophia Antipolis, France
- ²³INFN - Laboratori Nazionali del Sud (LNS), Via S. Sofia 62, 95123 Catania, Italy
- ²⁴Univ. Paris-Sud , 91405 Orsay Cedex, France
- ²⁵Kernfysisch Versneller Instituut (KVI), University of Groningen, Zernikelaan 25, 9747 AA Groningen, The Netherlands
- ²⁶Laboratoire de Physique Corpusculaire, Clermont Université, Université Blaise Pascal, CNRS/IN2P3, BP 10448, F-63000 Clermont-Ferrand, France
- ²⁷INFN - Sezione di Catania, Viale Andrea Doria 6, 95125 Catania, Italy
- ²⁸Royal Netherlands Institute for Sea Research (NIOZ), Landsdiep 4,1797 SZ 't Horntje (Texel), The Netherlands

- ²⁹Universiteit Utrecht, Faculteit Betawetenschappen, Princetonplein 5, 3584 CC Utrecht, The Netherlands
- ³⁰Dr. Remeis-Sternwarte and ECAP, Universität Erlangen-Nürnberg, Sternwartstr. 7, 96049 Bamberg, Germany
- ³¹Moscow State University, Skobeltsyn Institute of Nuclear Physics, Leninskie gory, 119991 Moscow, Russia
- ³²Dipartimento di Fisica ed Astronomia dell'Università, Viale Andrea Doria 6, 95125 Catania, Italy
- ³³Direction des Sciences de la Matière - Institut de recherche sur les lois fondamentales de l'Univers - Service de Physique des Particules, CEA Saclay, 91191 Gif-sur-Yvette Cedex, France
- ³⁴IPHC-Institut Pluridisciplinaire Hubert Curien - Université de Strasbourg et CNRS/IN2P3 23 rue du Loess, BP 28, 67037 Strasbourg Cedex 2, France
- ³⁵ITEP - Institute for Theoretical and Experimental Physics, B. Cheremushkinskaya 25, 117218 Moscow, Russia
- ³⁶Dipartimento di Fisica dell'Università, Via Dodecaneso 33, 16146 Genova, Italy
- ³⁷University Mohammed I, Laboratory of Physics of Matter and Radiations, B.P.717, Oujda 6000, Morocco
- ³⁸Institut für Theoretische Physik und Astrophysik, Universität Würzburg, Emil-Fischer-Str. 31, 97074 Würzburg, Germany
- ³⁹NASA, Goddard Space Flight Center, Greenbelt, MD 20771, USA
- ⁴⁰University of Maryland, Baltimore County, Baltimore, MD 21250, USA
- ⁴¹Catholic University of America, Washington, DC 20064, USA
- ⁴²Max-Planck-Institut für Radioastronomie, Auf dem Hgel 69, 53121 Bonn, Germany
- ⁴³Departament d'Astronomia i Astrofísica, Universitat de València, C/ Dr. Moliner 50, 46100 Burjassot, València, Spain
- ⁴⁴Observatori Astronòmic, Universitat de València, C/ Catedrático José Beltrán no. 2, 46980 Paterna, València, Spain
- ⁴⁵Departamento de Astronomía, Universidad de Concepción, Casilla 160, Chile
- ⁴⁶CSIRO Astronomy and Space Science, ATNF, PO Box 76 Epping, NSW 1710, Australia
- ⁴⁷Max-Planck-Institut für extraterrestrische Physik, Giessenbachstraße 1, 85741 Garching, Germany
- ⁴⁸Bundesamt für Kartographie und Geodäsie, 93444 Bad Kötzing, Germany
- ⁴⁹CSIRO Astronomy and Space Science, Canberra Deep Space Communications Complex, P.O. Box 1035, Tuggeranong, ACT 2901, Australia
- ⁵⁰School of Mathematics & Physics, University of Tasmania, Private Bag 37, Hobart, Tasmania 7001, Australia
- ⁵¹Hartebeesthoek Radio Astronomy Observatory, Krugersdorp, South Africa

February 2, 2015

Abstract

The source(s) of the neutrino excess reported by the IceCube Collaboration is unknown. The TANAMI Collaboration recently reported on the multiwavelength emission of six bright, variable blazars which are positionally coincident with two of the most energetic IceCube events. Such objects are prime candidates to be the source of the

*clancy.james@physik.uni-erlangen.de

†kadler@physik.uni-wuerzburg.de

highest-energy cosmic rays, and thus of associated neutrino emission. We present an analysis of neutrino emission from the six blazars using observations with the ANTARES neutrino telescope. The standard methods of the ANTARES candidate list search are applied to six years of data to search for an excess of muons — and hence their neutrino progenitors — from the directions of the six blazars described by the TANAMI Collaboration, and which are possibly associated with two IceCube events. Monte Carlo simulations of the detector response to both signal and background particle fluxes are used to estimate the sensitivity of this analysis for different possible source neutrino spectra. A maximum-likelihood approach, using the reconstructed energies and arrival directions of through-going muons, is used to identify events with properties consistent with a blazar origin. Both blazars predicted to be the most neutrino-bright in the TANAMI sample (1653–329 and 1714–336) have a signal flux fitted by the likelihood analysis corresponding to approximately one event. This observation is consistent with the blazar-origin hypothesis of the IceCube event IC14 for a broad range of blazar spectra, although an atmospheric origin cannot be excluded. No ANTARES events are observed from any of the other four blazars, including the three associated with IceCube event IC20. This excludes at a 90% confidence level the possibility that this event was produced by these blazars unless the neutrino spectrum is flatter than -2.4 .

1 Introduction

Since the initial report of the observation of two high-energy (\sim PeV) neutrino-induced cascades by the IceCube Collaboration [Aartsen et al., 2013], further observations using the high-energy starting-event (HESE) analysis have revealed an excess of events consistent with an isotropic, flavour-uniform flux of astrophysical neutrinos [IceCube Collaboration, 2013, Aartsen et al., 2014b,a]. The small number of excess events (37 total, with an estimated background of 15), and directional resolution of typically 10° or worse for cascades, makes it difficult to resolve potential features of this flux, such as a spectral downturn above PeV energies, a steeper spectral index, and/or a contribution from one or more point-like sources of neutrinos. Consequently, many suggestions for the nature and origin(s) of this flux have been put forward. Of particular note is the suggestion of a point-source near the Galactic Centre producing the observed excess in that region [Razzaque, 2013], a hypothesis already constrained by the ANTARES Collaboration [Adrián-Martínez et al., 2014a].

The TANAMI Collaboration has recently reported observations of six

bright, variable blazars (see Table 2) in positional coincidence with the range of possible arrival directions of the two PeV IceCube events IC14 and IC20 [Krauß et al., 2014]¹. Using a simple calculation based on the observed 1 keV to 10 GeV photon flux, the authors estimate that 1.9 ± 0.4 electron neutrino events at PeV energies would be expected in 662 days of IceCube data. This estimate compares well with the two observed events IC14 and IC20. Even taking this only as an order-of-magnitude indication of the expected event rate, a higher-resolution follow-up study of these objects is of great interest. Here, we present such an analysis using six years of data from the ANTARES neutrino telescope.

2 Target blazars and possible neutrino fluxes

The six blazars associated with the IC14 and IC20 fields by Krauß et al. [2014] are listed in Table 2. All exhibit prominent high-energy photon emission, and all but one are classified as flat-spectrum radio quasars (FSRQs) [Véron-Cetty and Véron, 2006]². The predictions of the expected number of detected electron neutrino events were made by assuming a neutrino energy $E_\nu = 1$ PeV and a flavour-uniform flux, with total energy flux equal to that in high-energy photons. Active galactic nuclei (AGN) of all classes have long been proposed as sites of hadronic interaction, and are potential sources of the highest-energy cosmic rays and, hence, neutrinos [Berezinskii and Smirnov, 1975, Hillas, 1984, Stecker and Salamon, 1996, Padovani and Resconi, 2014]. Predictions for the neutrino flux depend on the nature of the AGN considered, the cosmic-ray composition and flux, and the assumed densities of target hadronic matter and photon fields [Mannheim, 1995, Becker, 2008, Becker Tjus et al., 2014].

The emphasis on the two PeV events (IC14 and IC20; see Aartsen et al. [2014a] for a full list) comes from the fact that these two highest-energy events have only a negligible probability for an atmospheric origin. Moreover, Dermer et al. [2014] calculated that for typical FSRQs, the neutrino flux resulting from cosmic ray p - γ interactions should peak near a few PeV, with the primary target photon field being Lyman-series emission from the broad-line region. Other calculations of a neutrino flux from p - γ interactions

¹The paper was released before the third PeV event, IC35 (‘Big Bird’), was made public. A search for possible blazar associations with this event is in preparation by the TANAMI Collaboration.

²The exception is the source 1714–336, which has been classified as a BL Lac object. However, Krauß et al. [2014] find a prominent UV excess suggesting a possible reclassification as a FSRQ.

Source	Cat. Name	R.A. [$^{\circ}$]	Dec. [$^{\circ}$]	Class	z	F_{γ} [$\text{GeV cm}^{-2} \text{s}^{-1}$]	N_{ν_e}	IC
0235–618	PKS 0235–618	39.2218	–61.6043	Q	0.47 ^a	$(6.2^{+3.1}_{-3.1}) \times 10^{-8}$	$0.19^{+0.04}_{-0.04}$	20, 7
0302–623	PKS 0302–623	45.9610	–62.1904	Q	1.35 ^a	$(2.1^{+0.4}_{-0.4}) \times 10^{-8}$	$0.06^{+0.01}_{-0.01}$	20
0308–611	PKS 0308–611	47.4838	–60.9775	Q	1.48 ^a	$(4.7^{+1.8}_{-1.8}) \times 10^{-8}$	$0.14^{+0.05}_{-0.05}$	20
1653–329	Swift J1656.3–3302	254.0699	–33.0369	Q	2.40 ^b	$(2.8^{+0.3}_{-0.3}) \times 10^{-7}$	$0.86^{+0.10}_{-0.10}$	14, 2, 25
1714–336	TXS 1714–336	259.4001	–33.7024	B/Q	?	$(1.5^{+0.4}_{-0.4}) \times 10^{-7}$	$0.46^{+0.12}_{-0.12}$	14,2,25
1759–396	MRC 1759–396	270.6778	–39.6689	Q	1.32 ^c	$(7.5^{+1.9}_{-1.9}) \times 10^{-8}$	$0.23^{+0.50}_{-0.40}$	14, 2, 15, 25

Table 1: Basic data on the six blazars studied in this analysis. Columns: (1) IAU B1950 name; (2) Common catalog name; (3,4) J 2000 coordinates; (5) Classification: Q – Flat Spectrum Radio Quasar, B – BLLac object; (6) Redshift: ^a Healey et al. [2008], ^b Cutri et al. [2003], ^c Massaro et al. [2009]; (7) Total high-energy photon flux from Krauß et al. [2014]; (8) Estimated number N_{ν_e} of ν_e events in the IceCube 662-day analysis [IceCube Collaboration, 2013]; (9) IC gives the IceCube event IDs from Aartsen et al. [2014a] with which the blazars are positionally consistent within the angular error range from IceCube Collaboration [2013].

in blazars however have produced less-peaked spectra, e.g., that by Atoyan and Dermer [2001], who show that external radiation fields in 3C 279 can produce a flux of neutrinos slightly harder than E_ν^{-2} in the approximate range of 30 TeV–1 EeV.

Including p - p interactions can also lead to a neutrino flux at sub-PeV energies, which is expected to closely follow the cosmic-ray spectrum. In the case of a pure power-law proton spectrum $dN_p/dE_p \propto E_p^{-s_p}$, the pion, and hence — in a diffuse environment — neutrino spectrum reduces approximately to $dN_\nu/dE_\nu \propto E_\nu^{-s_\nu}$, with $s_\nu = \frac{4}{3}(s_p - \frac{1}{2})$ [Mannheim and Schlickeiser, 1994, Kelner et al., 2006]. Thus the simple Fermi acceleration model with $s_p = 2$ produces $s_\nu = 2$. More-detailed modelling of shock acceleration processes in AGN by Meli and Biermann [2012] has suggested that an initial accelerating shock in an AGN jet might produce an index of $s_p = 2.7$ ($s_\nu = 2.93$), while further shock acceleration leads to a flattening of the spectrum, producing $s_p = 2.4$ ($s_\nu = 2.53$) — which is nonetheless softer than the default $s_\nu = 2$ advocated by, e.g., Waxman and Bahcall [1999]. The incorporation of both p - p and p - γ interactions into full Monte Carlo calculations of particle interactions in a blazar environment confirms these expectations of a neutrino flux at sub-PeV energies [Szabo and Protheroe, 1994].

The IceCube observations allow for the possibility of a sub-PeV flux of neutrinos from the sample blazars, in that four other events are positionally associated with the blazar sample (see Table 2). This is also consistent with the prediction of two ν_e charged-current (CC) events, since the low flavour-dependence of the IceCube HESE effective area at the highest energies means an equal number of ν_μ and ν_τ events would be expected, but with a lower deposited energy. While these additional four events do not represent a significant excess above a diffuse background, the possibility that they may originate from the blazars in question should also be tested.

3 ANTARES candidate list search and expected sensitivity

ANTARES is an underwater neutrino telescope located in the Mediterranean Sea off the coast of Toulon, at $42^\circ 48'$ N, $6^\circ 10'$ E [Ageron et al., 2011]. Consisting of an array of photomultiplier tubes, it is designed to record the induced Cherenkov light from the passage of energetic charged particles to infer the interactions of neutrinos.

The ANTARES candidate list search (CLS) methodology is described in




Figure 1: (left) Relative exposures of the ANTARES CLS [Adrián-Martínez et al., 2012] to a flavour-uniform neutrino flux from the characteristic declinations of the six blazars of Table 2, and the Southern-Sky-average of the IceCube HESE analysis [Aartsen et al., 2014a] (exposures from IceCube Collaboration [2013]). The black dashed line, included for reference purposes, is proportional to E_ν . (right) Expected number of ANTARES events per detected IceCube event for power-law spectra (Eq. 1) as a function of the neutrino spectral index $-s_\nu$, calculated using the relative exposures.

Adrián-Martínez et al. [2012], with the latest results using six years of data (1338 days effective livetime) presented in Adrián-Martínez et al. [2014a]. The search uses only up-going muons (i.e., those originating from below the horizon), with cuts placed on the fit-quality of the muon track reconstruction and the estimated angular error. The long range of relativistic muons in seawater and the Earth’s crust extends the effective detection volume to well beyond the physical size of the detector, in contrast with a HESE-like analysis. The six-year sample consists of 5516 events, with an estimated atmospheric muon contamination of 10%, and an estimated median angular resolution of 0.38° . A maximum-likelihood method is then used to estimate the relative contributions of signal and background fluxes, based on both the reconstructed event arrival directions and the fitted number of photon hits (a robust proxy for energy). Note that this method results in a non-integer number of signal events N_{sig} being estimated, since the signal and background fluxes maximising the likelihood of a given observation can take any normalisation. Note also that it is optimised assuming an E_ν^{-2} source spectrum, and it is sensitive almost exclusively to muon neutrinos.

The ability of the ANTARES CLS to probe the PeV-neutrino blazar-origin hypotheses of Krauß et al. [2014] can be seen from Fig. 1, which compares the time-integrated, flavour-averaged exposures of the ANTARES CLS

(Adrián-Martínez et al. [2014a]; 1338 days, using one third of the effective area to muon neutrinos) at the characteristic declinations of the six blazars considered here, to that of the IceCube HESE analysis, averaged over the Southern Hemisphere (IceCube Collaboration [2013]; now updated to 998 days by Aartsen et al. [2014a], averaged over all three neutrino flavours). It can be seen that below approximately 100 TeV, ANTARES has a greater sensitivity to a neutrino flux from the six blazars at the given southern declinations than the recent IceCube HESE analysis. Due to the greater size of the IceCube detector, the effects of Earth absorption, and its sensitivity to all three neutrino flavours, the IceCube analysis has a significantly greater sensitivity at the highest energies. In particular, at an energy of 1 PeV, the exposure of IceCube to a flavour-uniform flux is approximately four times that of ANTARES. Therefore, the relative utility of an ANTARES analysis of the neutrino emission of these blazars will depend strongly on the energy spectrum of that flux.

The predictions for the number of IceCube-detected PeV neutrino events by Krauß et al. [2014], shown in Table 2, were based on equating the neutrino flux at 1 PeV to the integrated photon flux between 1 keV and 10 GeV. While — as discussed in Sect. 2 — the expected neutrino-flux shape is highly model-dependent, the prediction that the total neutrino energy flux F_ν ($\text{GeV cm}^{-2} \text{s}^{-1}$) is approximately equal to the total high-energy photon flux F_γ is relatively robust, at least when attributing this emission to a 100% hadronic origin. The black-dashed line in Fig. 1 is proportional to neutrino energy E_ν and normalised to the IceCube exposure at 1 PeV, i.e., it is a line of equal sensitivity to a neutrino flux F_ν . For constant F_ν , it is clear that the IceCube HESE analysis is most sensitive to a flux at a few hundred TeV, while the ANTARES CLS is most sensitive near 30 TeV.

A range of potential neutrino fluxes, $\Phi_\nu(E_\nu)$ (defined as the differential number-density flux dN_ν/dE_ν), can be characterised by generic power-law spectra of the form:

$$\Phi_\nu(E_\nu) = \Phi_0 \left(\frac{E_\nu}{1 \text{ GeV}} \right)^{-s_\nu} [\text{GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}]. \quad (1)$$

The relative numbers of events expected to be observed by ANTARES compared to IceCube for such spectra are shown in Fig. 1 (right). For a spectral index $-s_\nu < -2.2$, ANTARES is expected to observe more events, while IceCube would observe more for $-s_\nu > -2.2$.

For each spectral index $-s_\nu$ and source declination δ , the required neutrino flux $\Phi_\nu^*(E_\nu, \delta)$ expected to produce a single ANTARES event can be




Figure 2: Neutrino flux F_ν^* required to produce one neutrino event in ANTARES as a function of spectral index s_ν (Eq. 4). The corresponding energy ranges of integration E_{\min} and E_{\max} (Eq. 3) are shown as lower and upper shaded regions respectively — the shading covers the variation due to declination.

found from the following expression:

$$\int_0^\infty t_{\text{eff}} A_{\text{eff}}(E_\nu) \Phi_\nu^*(E_\nu, \delta) dE_\nu = 1, \quad (2)$$

where $A_{\text{eff}}(E_\nu, \delta)$ and t_{eff} are respectively the ANTARES effective area and observation time. While the total energy in such a flux is infinite, the energy over the sensitive range of ANTARES can be calculated by defining characteristic energies $E_{\min}(\delta, s_\nu)$ and $E_{\max}(\delta, s_\nu)$ such that:

$$\int_{E_{\min}}^{E_{\max}} t_{\text{eff}} A_{\text{eff}}(E_\nu, \delta) \Phi_\nu^*(E_\nu, \delta) dE_\nu = 0.9, \quad (3)$$

with 0.05 below E_{\min} and 0.05 above E_{\max} . The total neutrino energy flux $F_\nu^*(\delta, s_\nu)$ in the range $E_{\min} \leq E_\nu \leq E_{\max}$ required to produce one event can then be calculated from $\Phi_\nu^*(E_\nu, \delta)$ as:

$$F_\nu^*(\delta, s_\nu) = \frac{1}{0.9} \int_{E_{\min}}^{E_{\max}} \Phi_\nu^*(E_\nu, \delta) E_\nu dE_\nu \quad [\text{GeV cm}^{-2} \text{ s}^{-1}]. \quad (4)$$

$F_\nu^*(\delta, s_\nu)$ is plotted in Fig. 2, along with E_{\min} and E_{\max} . Comparing this with the total blazar photon flux calculated by Krauß et al. [2014] and reported in Table 2, it is clear that the detection of one or more neutrinos from power-law spectra in the range $-2.5 < -s_\nu < -1.5$ would be consistent with the observed source photon fluxes F_γ .

Having established a wide range of plausible neutrino flux scenarios, and the sensitivity of the ANTARES CLS to neutrino fluxes over a broad range of energies, we therefore perform the standard ANTARES CLS for an excess of neutrino emission from the blazars listed in Table 2.

4 Results and discussion

The results of the ANTARES analysis of the six blazars are given in Table 4. For four of the six targets, no source-like neutrinos were identified ($N_{\text{sig}} = 0$), allowing relatively strong upper limits to be placed on an E_ν^{-2} flux. The blazars 1653–329 and 1714–336 were each fitted as having approximately one nearby signal-like event, with N_{sig} of 1.1 and 0.9 respectively³. This observation is well within the expected background fluctuations however, with pre-trial p -values (probability of the likelihood procedure fitting a stronger signal flux to background-only data) of 0.10 and 0.04 respectively⁴. Nonetheless, it must be noted that these two blazars are the two with the highest predicted neutrino fluxes (see Table 2), and that from Fig. 1 (right), neutrino fluxes with spectral indices between -2.5 and -2.3 producing one IceCube event would be expected to produce between one and two ANTARES events. Therefore, when the calculation of Krauß et al. [2014] is extended to include power-law neutrino spectra, the result of the analysis is consistent with the sample blazars being neutrino sources with fluxes in proportion to their observed high-energy photon flux (F_γ in Table 2), even if the result is also consistent with background.

Limits at a 90% confidence level (C.L.), Φ_ν^{90} , on the spectra from Eq. 1 are generated from the ANTARES observations as a function of s_ν over the approximate predicted range (between 1.5 and 2.5), using the method of Neyman [1937]. These, all of which are upper limits, are given in Fig. 3 (left). Φ_ν^{90} is given at 100 TeV, because this is both the approximate energy at which the ANTARES and IceCube analyses have equal exposures, and where the flux limit is least sensitive to s_ν .

Constraints can be placed on a blazar origin of the IceCube events given in Table 2. The flux limits shown in Fig. 3 (left) correspond to a maximum expected number $N_{\nu,IC}^{90}$ of events observed by IceCube; where this number is less than the observed number of events, a blazar origin can be

³The maximum-likelihood procedure estimates N_{sig} as a continuous variable, as discussed in Sect. 3.

⁴The correct penalty factor for multiple trials is 61, including the six blazars considered here, and 55 trials from other analyses using the CLS [Adrián-Martínez et al., 2014a,b]

Source	N_{sig}	p	Limit $10^{-8} \text{ GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$	$N_{\nu, IC} = 1$	$N_{\nu, IC} = 2$	$N_{\nu, IC} = 3$	$N_{\nu, IC} = 4$
0235-618	0	1	1.3	-2.4	-2.1	-2.0	-1.9
0302-623	0	1	1.3	-2.4	-2.1	-2.0	-1.9
0308-611	0	1	1.3	-2.4	-2.1	-2.0	-1.9
1653-329	1.1	0.10	2.9	<-2.5	-2.5	-2.3	-2.2
1714-336	0.9	0.04	3.5	<-2.5	-2.5	-2.3	-2.2
1759-396	0	1	1.4	-2.4	-2.1	-2.0	-1.8

Table 2: ANTARES point-source analysis results. Columns: (1) IAU B 1950 name; (2) Number of fitted signal events; (3) pre-trial p -value; (4) 90% upper limit on Φ_0 for $-s_\nu = -2.0$, (5)-(8): minimum spectral indices $-s_\nu$ consistent at 90% C.L. with $N_{\nu, IC} = 1 \dots 4$ associated IceCube events.




Figure 3: (left) ANTARES 90% confidence limits on a flavour-uniform neutrino flux ($\Phi_\nu \equiv \Phi_{\nu_e} + \Phi_{\nu_\mu} + \Phi_{\nu_\tau} = 3\Phi_{\nu_\mu}$) from the six blazars as a function of spectral index s_ν (Eq. 1), and (right) corresponding limits on the expected number of IceCube events of blazar origin, using the exposures shown in Fig. 1 and the limiting fluxes. Since the limits from 0235–618, 0302–623, and 0308–611 are almost identical, and since no events were observed, the limits also apply to the summed flux from all three of these blazars, and hence only one line is shown, and labelled ‘IC20 TANAMI blazars’.

excluded at 90% C.L. This is shown in Fig. 3 (right). Any given number of IceCube events is therefore only consistent with a blazar origin for neutrino spectral indices flatter than certain value; minimum values of $-s_\nu$ are given for 1–4 events in Table 4, and should be compared to the possible associations in Table 2. For the IC14 field for instance, the possibility that blazar 1759–396 could be responsible for three or more associated IceCube events is excluded at 90% confidence for neutrino spectra steeper than -2.1 . For spectra steeper than -2.4 , we can exclude that 1759–396 is responsible for any IceCube events. The limits for 1653–329 and 1714–336 are weaker, due to a possible physical association with the two signal-like ANTARES events. Regardless of the association, we can rule out the possibility that the cluster IC14, IC2, and IC25 arose from a single considered blazar with a spectrum steeper than -2.4 . For the IC20 grouping, the non-observation of any event from the three candidate blazars means that the $\delta \approx -61^\circ$ limit applies both to the individual blazars, and the group as a whole. Therefore, ANTARES observations can rule out a neutrino spectrum steeper than -2.2 as being responsible for both IC20 and IC7, and a neutrino spectrum steeper than -2.4 being responsible for only one of them. That is, if IC20 does indeed originate from the three associated TANAMI blazars, the neutrino spectral index must be flatter than -2.4 .

5 Conclusion

We have tested the hypothesis of Krauß et al. [2014] that the first two PeV neutrino events observed by IceCube, IC14 and IC20, are of blazar origin, by performing a candidate list search (CLS) for an excess muon neutrino flux from the six suggested blazars using six years of ANTARES data. We are not able to either confirm or rule out a blazar origin of these events, although constraints have been placed on the range of source spectra which could have produced them, particularly in the case of IC20. While approximately two ANTARES events were fitted as being more signal-like than background-like by the maximum-likelihood analysis, such a result is completely within the expected background fluctuations, with pre-trial p -values of 10% and 4% for the blazars in question (1653–329 and 1714–336). It is interesting to note though that these two blazars were predicted by Krauß et al. [2014] to have the strongest neutrino flux, and that such a result is within expectations for the ANTARES event rate for an E_ν^{-2} to $E_\nu^{-2.3}$ neutrino spectrum given that IceCube observes two such events, and $E_\nu^{-2.3}$ to $E_\nu^{-2.5}$ for a single event of blazar origin. Given these considerations, the TANAMI candidate blazars should be included in all future analyses.

acknowledgement

The authors would like to thank A. Kappes for helpful discussions regarding the IceCube analysis. ANTARES authors acknowledge the financial support of the funding agencies: Centre National de la Recherche Scientifique (CNRS), Commissariat à l'énergie atomique et aux énergies alternatives (CEA), Commission Européenne (FEDER fund and Marie Curie Program), Région Alsace (contrat CPER), Région Provence-Alpes-Côte d'Azur, Département du Var and Ville de La Seyne-sur-Mer, France; Bundesministerium für Bildung und Forschung (BMBF), Germany; Istituto Nazionale di Fisica Nucleare (INFN), Italy; Stichting voor Fundamenteel Onderzoek der Materie (FOM), Nederlandse organisatie voor Wetenschappelijk Onderzoek (NWO), the Netherlands; Council of the President of the Russian Federation for young scientists and leading scientific schools supporting grants, Russia; National Authority for Scientific Research (ANCS), Romania; Ministerio de Ciencia e Innovación (MICINN), Prometeo of Generalitat Valenciana and MultiDark, Spain; Agence de l'Oriental and CNRST, Morocco. We also acknowledge the technical support of Ifremer, AIM and Foselev Marine for the sea operation and the CC-IN2P3 for the computing facilities. TANAMI authors acknowledge support and partial funding by the Deutsche Forschungsgemeinschaft grant WI

1860-10/1 (TANAMI) and GRK 1147, Deutsches Zentrum für Luft- und Raumfahrt grant 50 OR 1311/50 OR 1303/50 OR 1401, the Spanish MINECO project AYA2012-38491-C02-01, the Generalitat Valenciana project PROMETEOII/2014/057, the COST MP0905 action “Black Holes in a Violent Universe” and the Helmholtz Alliance for Astroparticle Physics (HAP).

References

- M. G. Aartsen, R. Abbasi, Y. Abdou, M. Ackermann, J. Adams, J. A. Aguilar, M. Ahlers, D. Altmann, J. Auffenberg, X. Bai, and et al. First Observation of PeV-Energy Neutrinos with IceCube. *Physical Review Letters*, 111(2):021103, July 2013. doi: 10.1103/PhysRevLett.111.021103.
- M. G. Aartsen, M. Ackermann, J. Adams, J. A. Aguilar, M. Ahlers, M. Ahrens, D. Altmann, T. Anderson, C. Argüelles, T. C. Arlen, and et al. Observation of High-Energy Astrophysical Neutrinos in Three Years of IceCube Data. *Physical Review Letters*, 113(10):101101, Sept. 2014a. doi: 10.1103/PhysRevLett.113.101101.
- M. G. Aartsen, M. Ackermann, J. Adams, J. A. Aguilar, M. Ahlers, M. Ahrens, D. Altmann, T. Anderson, C. Argüelles, T. C. Arlen, and et al. Atmospheric and Astrophysical Neutrinos above 1 TeV Interacting in IceCube. *ArXiv e-prints*, Oct. 2014b. arXiv:1410.1749.
- S. Adrián-Martínez et al. Search for Cosmic Neutrino Point Sources with Four Years of Data from the ANTARES Telescope. *ApJ*, 760:53, Nov. 2012. doi: 10.1088/0004-637X/760/1/53.
- S. Adrián-Martínez, et al. Searches for Point-like and Extended Neutrino Sources Close to the Galactic Center Using the ANTARES Neutrino Telescope. *ApJ*, 786:L5, May 2014a. doi: 10.1088/2041-8205/786/1/L5.
- S. Adrián-Martínez, et al. Constraining the neutrino emission of gravitationally lensed Flat-Spectrum Radio Quasars with ANTARES data. *ArXiv e-prints*, July 2014b. arXiv:1407.8525.
- M. Ageron, J. A. Aguilar, I. Al Samarai, A. Albert, F. Ameli, M. André, M. Anghinolfi, G. Anton, S. Anvar, M. Ardid, and et al. ANTARES: The first undersea neutrino telescope. *Nuclear Instruments and Methods in Physics Research A*, 656:11–38, Nov. 2011. doi: 10.1016/j.nima.2011.06.103.

- A. Atoyan and C. D. Dermer. High-Energy Neutrinos from Photomeson Processes in Blazars. *Physical Review Letters*, 87(22):221102, Nov. 2001. doi: 10.1103/PhysRevLett.87.221102.
- J. K. Becker. High-energy neutrinos in the context of multimessenger astrophysics. *Phys. Rep.*, 458:173–246, Mar. 2008. doi: 10.1016/j.physrep.2007.10.006.
- J. Becker Tjus, B. Eichmann, F. Halzen, A. Kheirandish, and S. M. Saba. High-energy neutrinos from radio galaxies. *Phys. Rev. D*, 89(12):123005, June 2014. doi: 10.1103/PhysRevD.89.123005.
- V. S. Berezinskii and A. I. Smirnov. Cosmic neutrinos of ultra-high energies and detection possibility. *Ap&SS*, 32:461–482, Feb. 1975. doi: 10.1007/BF00643157.
- R. M. Cutri et al. VizieR Online Data Catalog: 2MASS All-Sky Catalog of Point Sources (Cutri+ 2003). *VizieR Online Data Catalog*, 2246:0, Mar. 2003.
- C. D. Dermer, K. Murase, and Y. Inoue. Photopion production in black-hole jets and flat-spectrum radio quasars as PeV neutrino sources. *Journal of High Energy Astrophysics*, 3:29–40, Sept. 2014. doi: 10.1016/j.jheap.2014.09.001.
- S. E. Healey, R. W. Romani, G. Cotter, P. F. Michelson, E. F. Schlafly, A. C. S. Readhead, P. Giommi, S. Chaty, I. A. Grenier, and L. C. Weintraub. CGRaBS: An All-Sky Survey of Gamma-Ray Blazar Candidates. *ApJS*, 175:97–104, Mar. 2008. doi: 10.1086/523302.
- A. M. Hillas. The Origin of Ultra-High-Energy Cosmic Rays. *ARA&A*, 22: 425–444, 1984. doi: 10.1146/annurev.aa.22.090184.002233.
- IceCube Collaboration. Evidence for High-Energy Extraterrestrial Neutrinos at the IceCube Detector. *Science*, 342:1242856, Nov. 2013. doi: 10.1126/science.1242856.
- S. R. Kelner, F. A. Aharonian, and V. V. Bugayov. Energy spectra of gamma rays, electrons, and neutrinos produced at proton-proton interactions in the very high energy regime. *Phys. Rev. D*, 74:034018, Aug 2006. doi: 10.1103/PhysRevD.74.034018. URL <http://link.aps.org/doi/10.1103/PhysRevD.74.034018>.

- F. Krauß , et al. TANAMI blazars in the IceCube PeV-neutrino fields. *A&A*, 566:L7, June 2014. doi: 10.1051/0004-6361/201424219.
- K. Mannheim. High-energy neutrinos from extragalactic jets. *Astroparticle Physics*, 3:295–302, May 1995. doi: 10.1016/0927-6505(94)00044-4.
- K. Mannheim and R. Schlickeiser. Interactions of cosmic ray nuclei. *A&A*, 286:983–996, June 1994.
- E. Massaro, P. Giommi, C. Leto, P. Marchegiani, A. Maselli, M. Perri, S. Piranomonte, and S. Scavi. Roma-BZCAT: a multifrequency catalogue of blazars. *A&A*, 495:691–696, Feb. 2009. doi: 10.1051/0004-6361:200810161.
- A. Meli and P. L. Biermann. UHECRs and multiple shock acceleration in Active Galactic Nuclei Jets. *Journal of Physics Conference Series*, 375(5):052016, July 2012. doi: 10.1088/1742-6596/375/1/052016.
- J. Neyman. Outline of a Theory of Statistical Estimation Based on the Classical Theory of Probability. *Royal Society of London Philosophical Transactions Series A*, 236:333–380, Aug. 1937. doi: 10.1098/rsta.1937.0005.
- P. Padovani and E. Resconi. Are both BL Lacs and pulsar wind nebulae the astrophysical counterparts of IceCube neutrino events? *MNRAS*, 443:474–484, Sept. 2014. doi: 10.1093/mnras/stu1166.
- S. Razzaque. Galactic Center origin of a subset of IceCube neutrino events. *Phys. Rev. D*, 88(8):081302, Oct. 2013. doi: 10.1103/PhysRevD.88.081302.
- F. W. Stecker and M. H. Salamon. High Energy Neutrinos from Quasars. *Space Sci. Rev.*, 75:341–355, Jan. 1996. doi: 10.1007/BF00195044.
- A. P. Szabo and R. J. Protheroe. Implications of particle acceleration in active galactic nuclei for cosmic rays and high energy neutrino astronomy. *Astroparticle Physics*, 2:375–392, Oct. 1994. doi: 10.1016/0927-6505(94)90027-2.
- M.-P. Véron-Cetty and P. Véron. A catalogue of quasars and active nuclei: 12th edition. *A&A*, 455:773–777, Aug. 2006. doi: 10.1051/0004-6361:20065177.

E. Waxman and J. Bahcall. High energy neutrinos from astrophysical sources: An upper bound. *Phys. Rev. D*, 59(2):023002, Jan. 1999. doi: 10.1103/PhysRevD.59.023002.