

## 3-permutable subgroups of finite groups

A. A. Heliel · A. Ballester-Bolinches · R.  
Esteban-Romero · M. O. Almistady

**Abstract** Let  $\mathfrak{S}$  be a complete set of Sylow subgroups of a finite group  $G$ , that is, a set composed of a Sylow  $p$ -subgroup of  $G$  for each  $p$  dividing the order of  $G$ . A subgroup  $H$  of  $G$  is called  $\mathfrak{S}$ -permutable if  $H$  permutes with all members of  $\mathfrak{S}$ . The main goal of this paper is to study the embedding of the  $\mathfrak{S}$ -permutable subgroups and the influence of  $\mathfrak{S}$ -permutability on the group structure.

**Keywords** finite group,  $p$ -soluble group,  $p$ -supersoluble group,  $\mathfrak{S}$ -permutable subgroup, subnormal subgroup

**Mathematics Subject Classification (2000)** 20D10 · 20D20 · 20D35 · 20D40

### 1 Introduction and statements of results

All groups in this paper will be finite.

---

A. A. Heliel

Department of Mathematics, Faculty of Science 80203, King Abdulaziz University, Jeddah 21589, Saudi Arabia, and Department of Mathematics, Faculty of Science 62511, Beni-Suef University Beni-Suef, Egypt

E-mail: heliel9@yahoo.com

A. Ballester-Bolinches

Departament d'Àlgebra, Universitat de València; Dr. Moliner, 50; 46100, Burjassot, València, Spain

E-mail: Adolfo.Ballester@uv.es

R. Esteban-Romero

Institut Universitari de Matemàtica Pura i Aplicada, Universitat Politècnica de València; Camí de Vera, s/n; 46022 València, Spain

E-mail: resteban@mat.upv.es

*Present adress:* Departament d'Àlgebra, Universitat de València; Dr. Moliner, 50; 46100, Burjassot, València, Spain

E-mail: Ramon.Esteban@uv.es

M. O. Almistady

Department of Mathematics, Faculty of Science 80203, King Abdulaziz University; Jeddah 21589, Saudi Arabia

A subgroup  $H$  of a group  $G$  is said to *permute* with a subgroup  $K$  of  $G$  if  $HK$  is a subgroup of  $G$ .  $H$  is said to be *S-permutable* in  $G$  if  $H$  permutes with all Sylow subgroups of  $G$ . This property extends normality and permutability and was introduced by Kegel in [11]. In this paper, he proved some interesting properties which turn out to be useful in establishing results concerning the group structure. In particular, it is proved there that every S-permutable subgroup must be subnormal ([11, Satz 1]).

On the other hand, we say that a set  $\mathfrak{S}$  of Sylow subgroups of a group  $G$  is a *complete set of Sylow subgroups of  $G$*  if  $\mathfrak{S}$  contains exactly one Sylow subgroup for each prime dividing the order of  $G$ ;  $\mathfrak{S}$  is called a *Sylow basis of  $G$*  if the Sylow subgroups in  $\mathfrak{S}$  are pairwise permutable. Sylow basis were introduced and studied by Hall in [7]. The results of this paper show that the existence and conjugacy of Sylow bases is a characteristic property of soluble groups.

In [1], Asaad and Heliel introduced and studied the notion of a  $\mathfrak{S}$ -permutable subgroup, where  $\mathfrak{S}$  is a complete set of Sylow subgroups of a group  $G$ . A subgroup of  $G$  is called  *$\mathfrak{S}$ -permutable* if it permutes with every member of a complete set  $\mathfrak{S}$  of Sylow subgroups of  $G$ . It is clear that S-permutability implies  $\mathfrak{S}$ -permutability but the converse does not hold in general. In fact,  $\mathfrak{S}$ -permutable subgroups are not subnormal in general, and subnormal  $\mathfrak{S}$ -permutable subgroups are not S-permutable either as the following example shows:

*Example 1* Let  $E = \langle x, y \rangle$  be an extraspecial group of order 27 and exponent 3. Let  $a$  be an automorphism of order 2 of  $E$  given by  $x^a = x^{-1}$ ,  $y^a = y^{-1}$ . Let  $G = E \rtimes \langle a \rangle$  be the corresponding semidirect product. Then  $\mathfrak{S} = \{E, \langle a \rangle\}$  is a complete set of Sylow subgroups of  $G$ . The subgroup  $H = \langle x \rangle$  is  $\mathfrak{S}$ -permutable, but it does not permute with the Sylow 2-subgroup  $\langle a \rangle$ . Therefore,  $H$  is not S-permutable. However,  $H$  is a subnormal subgroup of  $G$ .

Throughout the first part of our paper, proving important properties of S-permutable type of the subnormal  $\mathfrak{S}$ -permutable subgroups has been our main focus.

The embedding of S-permutable subgroups was studied by Kegel [11, Satz 1] and Deskins [4, Theorem 1] (see also [3, Theorem 1.2.14]). They proved that if  $A$  is an S-permutable subgroup of  $G$ , then  $\langle A^G \rangle / \text{Core}_G(A)$  is nilpotent. Our first main result shows how a subnormal  $\mathfrak{S}$ -permutable subgroup is embedded in the group.

**Theorem 1** *Let  $\mathfrak{S}$  be a complete set of Sylow subgroups of a group  $G$ . Let  $A$  be a subnormal  $\mathfrak{S}$ -permutable subgroup of  $G$ . Then  $\langle A^G \rangle / \text{Core}_G(A)$  is soluble. If, in addition,  $\mathfrak{S}$  is a Sylow basis of  $G$ , then  $\langle A^G \rangle / \text{Core}_G(A)$  is nilpotent.*

The alternating group of degree 6 is a non-subnormal  $\mathfrak{S}$ -permutable subgroup of the alternating group of degree 7 which is not soluble. Moreover, every core-free maximal subgroup of a soluble primitive group is  $\mathfrak{S}$ -permutable. Therefore subnormality is necessary in the above theorem.

A classical result of Kegel ([11, Satz 2], see also [3, Theorem 1.2.19]) shows that the set of all S-permutable subgroups of a group  $G$  is a sublattice of the subgroup lattice of  $G$ . Kegel's result also holds for subnormal  $\mathfrak{S}$ -permutable subgroups. It is consequence of the following theorem.

**Theorem 2** *Let  $p$  be a prime and  $U$  and  $V$  subgroups of a group  $G$ . If  $U$  and  $V$  permute with a Sylow  $p$ -subgroup  $G_p$  of  $G$  and  $U$  is subnormal in  $G$ , then  $U \cap V$  permutes with  $G_p$ .*

The hypothesis of the subnormality of  $U$  is necessary in the above theorem, even for soluble groups, as an example of Doerk [5, Beispiel 1] shows.

**Corollary 1** *Let  $\mathfrak{S}$  be a complete set of Sylow subgroups of a group  $G$ . Then the set of all subnormal  $\mathfrak{S}$ -permutable subgroups of a group  $G$  is a sublattice of the lattice of all subgroups of  $G$ .*

If  $\mathfrak{S}$  is a Sylow basis of a group  $G$ , the set of all  $\mathfrak{S}$ -permutable subgroups is a sublattice of the subgroup lattice of  $G$  ([6, Chapter I, Theorem 4.29]). However, we do not know whether the set of all  $\mathfrak{S}$ -permutable subgroups (not necessarily subnormal) of a group  $G$  is a sublattice of the lattice of all subgroups of  $G$ .

There are several papers in the literature where global information about a group is obtained by assuming that some distinguished subgroups are  $\mathfrak{S}$ -permutable ([1, 8, 9, 12, 13, 14, 15, 17]). The second part of the paper concerns situations in this spirit, but we require only that some  $p$ -subgroups, for a given prime  $p$ , have the required property.

In order to state our results in this direction, we recall that a group is said to be  $p$ -supersoluble if it is  $p$ -soluble and every  $p$ -chief factor has order  $p$ , where  $p$  is a prime that we hold fixed.

In the sequel,  $\mathfrak{S} = \{G_q \mid q \in \pi(G)\}$  will denote a complete set of Sylow subgroups of a group  $G$ , where  $G_q$  is a Sylow  $q$ -subgroup of  $G$ .

Asaad and Heliel [1, Theorem 3.1] showed that if all maximal subgroups of the Sylow subgroups in  $\mathfrak{S}$  are  $\mathfrak{S}$ -permutable, then  $G$  is supersoluble. A local approach to this theorem is the following.

**Theorem 3** *Let  $G$  be a group. Assume that all maximal subgroups of  $G_p \in \mathfrak{S}$  are  $\mathfrak{S}$ -permutable. Then either  $G_p$  is cyclic or  $G$  is  $p$ -supersoluble.*

If  $p$  is the smallest prime dividing the order of  $G$ , and the Sylow  $p$ -subgroups are cyclic, then  $G$  is  $p$ -nilpotent by [10, IV, Satz 2.8]. Furthermore, if  $G$  is  $p$ -supersoluble, then  $G$  every  $p$ -chief factor is central and so  $G$  is  $p$ -nilpotent. Therefore we have:

**Corollary 2** ([14, Theorem 3.1]) *If  $p$  is the smallest prime dividing the order of a group  $G$  and the maximal subgroups of  $G_p \in \mathfrak{S}$  are  $\mathfrak{S}$ -permutable, then  $G$  is  $p$ -nilpotent.*

**Corollary 3** ([1, Theorem 3.1]) *Assume that  $G$  is a group whose maximal subgroups of the Sylow subgroups in  $\mathfrak{S}$  are  $\mathfrak{S}$ -permutable. Then  $G$  is supersoluble.*

The next natural step in our analysis to consider the structural impact of the  $\mathfrak{S}$ -permutability of the second maximal subgroups of the Sylow  $p$ -subgroup in  $\mathfrak{S}$ .

Suppose that every 2-maximal subgroup of  $G_p \in \mathfrak{S}$  is  $\mathfrak{S}$ -permutable and that  $G_p$  does not have cyclic maximal subgroups. Then every maximal subgroup of  $G_p$  is  $\mathfrak{S}$ -permutable and  $G_p$  is not cyclic. By Theorem 3,  $G$  is  $p$ -supersoluble. Therefore we have:

**Corollary 4** *Let  $G$  be a group. Suppose that all 2-maximal subgroups of  $G_p \in \mathfrak{3}$  are  $\mathfrak{3}$ -permutable. Either  $G_p$  has a cyclic maximal subgroup or  $G$  is  $p$ -supersoluble.*

In [14, Theorem 3.3], the authors proved the following result:

**Theorem 4 ([14, Theorem 3.3])** *Assume that  $p$  is the smallest prime dividing the order of a group  $G$ . Suppose that all 2-maximal subgroups of  $G_p \in \mathfrak{3}$  are  $\mathfrak{3}$ -permutable. If  $G$  is  $A_4$ -free, then  $G$  is  $p$ -nilpotent.*

Our goal in the sequel is to present an improvement of this theorem.

According to Corollary 4, if all 2-maximal subgroups of  $G_p \in \mathfrak{3}$  are  $\mathfrak{3}$ -permutable, then either  $G_p$  has a cyclic maximal subgroup or  $G$  is  $p$ -supersoluble. Furthermore, a  $p$ -supersoluble group  $G$  is  $p$ -nilpotent provided that  $p$  is the smallest prime dividing the order of  $G$ . Hence we only must consider the case when  $G_p$  has a cyclic maximal subgroup. We prove:

**Theorem 5** *Assume that  $p$  is the smallest prime dividing the order of a group  $G$ . Suppose that all 2-maximal subgroups of  $G_p \in \mathfrak{3}$  are  $\mathfrak{3}$ -permutable. If  $G_p$  has a cyclic maximal subgroup, then  $G$  is  $p$ -soluble.*

A key fact for the proof of Theorem 5 is that  $G$  cannot be non-abelian simple. This was established in Step 3 of the proof of [14, Theorem 3.3].

**Theorem 6** *Assume that  $p$  is the smallest prime dividing the order of a group  $G$ . If every 2-maximal subgroup of  $G_p \in \mathfrak{3}$  is  $\mathfrak{3}$ -permutable, then either  $G$  is  $p$ -nilpotent or  $G$  has an epimorphic image isomorphic to  $\Sigma_4$ .*

In [12, Theorem 3.3], the authors proved that if  $p$  is the smallest prime dividing the order of a group of  $G$  and every cyclic subgroup of  $G_p$  with order  $p$  or order 4 (if  $p = 2$ ) is  $\mathfrak{3}$ -permutable in  $G$ , then  $G$  is  $p$ -nilpotent.

Our last results concern the  $\mathfrak{3}$ -permutability of the minimal subgroups of the Sylow  $p$ -subgroup in  $\mathfrak{3}$  and include the above result as a particular case.

**Theorem 7** *Let  $G$  be a  $p$ -soluble group such that every cyclic subgroup of  $G_p$  with order  $p$  or order 4 (if  $p = 2$ ) is  $\mathfrak{3}$ -permutable in  $G$ . Then  $G$  is  $p$ -supersoluble.*

**Theorem 8** *Let  $G$  be a group such that every cyclic subgroup of  $G_p$  with order  $p$  or order 4 (if  $p = 2$ ) is  $\mathfrak{3}$ -permutable in  $G$ . Either  $G_p$  has order  $p$  or  $G$  is  $p$ -soluble.*

**Corollary 5** *Let  $G$  be a group such that every cyclic subgroup of  $G_p$  with order  $p$  or order 4 (if  $p = 2$ ) is  $\mathfrak{3}$ -permutable in  $G$ . Then either  $G_p$  has order  $p$  or  $G$  is  $p$ -supersoluble.*

**Corollary 6 ([12, Theorem 3.3])** *If  $p$  is the smallest prime dividing the order of  $G$  and every cyclic subgroup of  $G_p$  with order  $p$  or order 4 (if  $p = 2$ ) is  $\mathfrak{3}$ -permutable in  $G$ , then  $G$  is  $p$ -nilpotent.*

## 2 Preliminaries

Suppose that  $G$  is a group and  $N$  a normal subgroup of  $G$ . Following [1], we write  $\mathfrak{Z}N = \{G_q N : G_q \in \mathfrak{Z}\}$ ,  $\mathfrak{Z}N/N = \{G_q N/N : G_q \in \mathfrak{Z}\}$ , and  $\mathfrak{Z} \cap X = \{G_q \cap X : G_q \in \mathfrak{Z}\}$  for all subgroups  $X$  of  $G$ .

**Lemma 1** ([1, Lemma 2.1]) *Let  $G$  be a group and  $N$  a normal subgroup of  $G$ .*

1.  $\mathfrak{Z} \cap N$  and  $\mathfrak{Z}N/N$  are complete sets of Sylow subgroups of  $N$  and  $G/N$ , respectively.
2. If  $U$  is a  $\mathfrak{Z}$ -permutable subgroup of  $G$ , then  $UN/N$  is  $\mathfrak{Z}N/N$ -permutable. If  $U$  is contained in  $N$ , then  $U$  is  $\mathfrak{Z} \cap N$ -permutable.

The following well-known fact, which follows from the repeated application of [10, Kapitel I, Hilfssatz 7.7a)], will be used in this paper without further notice.

**Lemma 2** *Let  $S$  be a subnormal subgroup of a group  $G$  and let  $Q$  be a Sylow  $q$ -subgroup of  $G$ , where  $q$  is a prime. Then  $Q \cap S$  is a Sylow  $q$ -subgroup of  $S$ .*

The following result, due to Vdovin, turns out to be crucial in the proofs of some of our results.

**Theorem 9** *If, for every prime  $q \neq p$ ,  $G$  possesses a Hall  $\{p, q\}$ -subgroup, then  $G$  is  $p$ -soluble.*

The above theorem is a consequence of the following lemma, whose proof requires a bit of notation.

Let  $q$  be a natural number, and  $r$  an odd prime such that  $\gcd(q, r) = 1$ . Let  $e(q, r)$  denote the multiplicative order of  $q$  modulo  $r$ , that is, the least natural number  $t$  with  $q^t \equiv 1 \pmod{r}$ . For an odd  $q$ , we set  $e(q, 2) = 1$  if  $q \equiv 1 \pmod{4}$  and  $e(q, 2) = 2$  otherwise.

**Lemma 3** *Let  $r$  be a prime. Then, for every simple group  $S$  with  $r \in \pi(S)$ , there exists  $s \in \pi(S)$  such that  $S$  does not possess a Hall  $\{r, s\}$ -subgroup.*

*Proof* Suppose, by contradiction, that there exists a finite simple group  $S$  and a prime  $r \in \pi(S)$  such that, for every  $s \in \pi(S)$ ,  $S$  possesses a Hall  $\{r, s\}$ -subgroup  $H$ . Burnside's  $p^a q^b$ -theorem implies that  $H$  is soluble. We proceed case by case.

Assume first that  $S$  is an alternating group  $A_n$  of degree  $n \geq 5$ . If  $r \neq 2$ , then [16, Table 2] implies that for every odd  $s \in \pi(G) \setminus \{p\}$ , we have that  $S$  does not possess a Hall  $\{r, s\}$ -subgroup. If  $r = 2$ , then [16, Table 2] implies that  $S$  does not have a Hall  $\{2, 5\}$ -subgroup.

Now assume that  $S$  is sporadic. Then the claim follows from [16, Tables 3 and 4].

Finally, assume that  $S$  is a finite simple group of Lie type over a field of characteristic  $p$  and order  $q$ . If  $r = p$ , then [16, Theorem 8.3] implies that every Hall  $\pi$ -subgroup of  $S$  with  $r \in \pi$  is contained in a Borel subgroup  $B$  or is parabolic. Since  $B$  is a proper subgroup of  $S$ , there exists  $s \in \pi(|S : B|)$ , and so  $B$  cannot contain a Hall  $\{r, s\}$ -subgroup of  $S$ . Therefore a Hall  $\{r, s\}$ -subgroup  $H$  of  $S$  is parabolic. Theorems 8.5, 8.6, and 8.7 and Table 6 from [16] imply that in this case  $\{r, s\} = \{2, 3\}$

and  $S \in \{\mathrm{SL}_3(2), \mathrm{SL}_3(3), \mathrm{SL}_4(2), \mathrm{SL}_5(2)\}$ . In all these cases, there is no Hall  $\{r, s\}$ -subgroup for  $s \in \pi((q^n - 1)/(q - 1))$ , which is always contained in  $\pi' \cap \pi(G)$  if  $G$  has a proper Hall  $\pi$ -subgroup with  $|\pi| \geq 2$ .

Assume that  $r \neq p$  is that  $r$  is odd. If  $S$  is an exceptional group of Lie type and  $S$  is neither a Suzuki nor a Ree group, then [16, Table 7] implies that  $S \in E_{\{r,s\}}$  if and only if  $e(q, r) = e(q, s)$ . Now by the decomposition of  $|S|$  as a product of polynomials in  $q$ , there exists an odd  $s \in \pi(S)$  with  $e(q, r) \neq e(q, s)$ , and so  $S$  does not have a Hall  $\{r, s\}$ -subgroup. If  $S$  is either a Suzuki or a Ree group, then the claim follows immediately from [16, Table 8]. If  $S$  is a classical group of Lie type, then [16, Table 7] implies that  $S$  has a Hall  $\{r, s\}$ -subgroup only if either  $e(q, r) = e(q, s)$  or  $e(q, s) = b(r)$ , where  $b(r) \in \{1, r\}$  if  $G = \mathrm{PSL}_n(q)$ ,  $b(r) \in \{2, 2r\}$  if  $G = \mathrm{PSU}_n(q)$ ,  $b(r) = 2e(q, r)$  if  $e(q, r)$  is odd and  $G = {}^2D_n(q)$ ,  $b(r) = e(q, r)/2$  if  $e(q, r)$  is even, 4 does not divide  $e(q, r)$  and  $G = {}^2D_n(q)$ . In particular, if  $e(q, s) \neq e(q, r)$ , then  $e(q, s)$  can take at most two values. If the rank of  $S$  is at least 2, then  $|\{e(q, s) \mid s \in \pi(S) \setminus \{p\}\}| \geq 3$ , and so there exists  $s$  such that  $e(q, r) \notin \{e(q, r), b(r)\}$  and therefore  $S \notin E_{\{r,s\}}$ . If the rank of  $S$  is less than 2, then  $S \cong \mathrm{PSL}_2(q)$ . If  $S = \mathrm{PSL}_2(q)$  and  $e(q, r) = 1$ , then there exists  $s \in \pi(S)$  with  $e(q, s) = 2$  and [16, Tables 7 and 10] imply that  $S \notin E_{\{r,s\}}$ . If  $S \cong \mathrm{PSL}_2(q)$  and  $e(q, r) = 2$ , then  $S \notin E_{\{r,p\}}$ .

Finally, assume that  $r = 2$  and  $r \neq p$ . If  $S \neq \mathrm{SL}_3(3)$ , then  $S \notin E_{\{2,p\}}$  by the arguments presented for the case  $r = p$ . If  $S = \mathrm{SL}_3(3)$ , then  $S \notin E_{\{2,13\}}$ .

**Corollary 7** *Let  $G$  be a group and  $p \in \pi(G)$ . Assume that*

1. *all maximal subgroups of  $G_p \in \mathfrak{F}$  are  $\mathfrak{F}$ -permutable and  $G_p$  is not cyclic, or*
2. *all 2-maximal subgroups of  $G_p \in \mathfrak{F}$  are  $\mathfrak{F}$ -permutable and  $G_p$  has no cyclic maximal subgroups.*

*Then  $G$  is  $p$ -soluble.*

*Proof* Assume that 1 holds. Then  $G_p$  possesses two maximal subgroups  $M_1$  and  $M_2$ , both  $\mathfrak{F}$ -permutable. Then  $M_1M_2 = G_p$  is  $\mathfrak{F}$ -permutable. This implies that  $G_pG_q$  is a Hall  $\{p, q\}$ -subgroup of  $G$  for each  $q \neq p$ . By Theorem 9,  $G$  is  $p$ -soluble.

Assume that 2 holds. Let  $M_1$  be a maximal subgroup of  $G_p$ . Since  $M_1$  is not cyclic,  $M_1$  possesses two maximal subgroups  $M_{11}$  and  $M_{12}$ . Since both of them are  $\mathfrak{F}$ -permutable,  $M_1 = M_{11}M_{12}$  is also  $\mathfrak{F}$ -permutable. Hence 1 holds and  $G$  is  $p$ -soluble.

### 3 Proofs of the main results

*Proof (of Theorem 1)* We prove that  $A/\mathrm{Core}_G(A)$  is soluble by induction on the order of  $G$ . Since  $A/\mathrm{Core}_G(A)$  is  $(\mathfrak{F}\mathrm{Core}_G(A)/\mathrm{Core}_G(A))$ -permutable in  $G/\mathrm{Core}_G(A)$  by Lemma 1, we can assume that  $\mathrm{Core}_G(A) = 1$ . Let  $r$  be a prime dividing  $|G|$  and let  $R$  be the Sylow  $r$ -subgroup of  $G$  in  $\mathfrak{F}$ . Consider  $X = AR$ . Let  $q$  be a prime different from  $r$  and let  $G_q$  be the Sylow  $q$ -subgroup of  $G$  in  $\mathfrak{F}$ . Since  $A$  is subnormal in  $G$ ,  $G_q \cap A$  is a Sylow  $q$ -subgroup of  $A$ . Moreover,  $A$  is  $(\mathfrak{F} \cap X)$ -permutable, because  $\mathfrak{F} \cap X = \{R\} \cup \{G_q \cap A \mid q \neq r\}$ . Moreover  $A$  is subnormal in  $X$ . Assume that  $X$  is a proper subgroup of  $G$ . By induction, the soluble residual  $A^\mathfrak{S}$  of  $A$  is contained in  $\mathrm{Core}_X(A) = \mathrm{Core}_R(A)$ . Consequently,  $A^\mathfrak{S} = (A^\mathfrak{S})^\mathfrak{S} \leq \mathrm{Core}_R(A)^\mathfrak{S} \leq A^\mathfrak{S}$ . It follows

that  $A^\mathfrak{S} = \text{Core}_R(A)^\mathfrak{S}$  is a normal subgroup of  $X$ . In particular,  $R \leq N_G(A^\mathfrak{S})$ . Suppose that for every Sylow subgroup  $R$  of  $G$  in  $\mathfrak{S}$ ,  $AR$  is a proper subgroup of  $G$ . It follows that  $R \leq N_G(A^\mathfrak{S})$  for each  $R \in \mathfrak{S}$ . Hence  $A^\mathfrak{S}$  is a normal subgroup of  $G$ . Thus  $A^\mathfrak{S} \leq \text{Core}_G(A) = 1$ . Consequently  $A$  is soluble, as wanted.

Therefore there exists a prime  $r$  and a Sylow  $r$ -subgroup  $R$  of  $G$  in  $\mathfrak{S}$  such that  $G = AR$ . Let  $q$  be a prime different from  $r$  and let  $Q$  be a Sylow  $q$ -subgroup of  $G$ . The subnormality of  $A$  implies that  $Q \cap A$  is a Sylow  $q$ -subgroup of  $A$ . By order considerations,  $Q \cap A = Q$  and so  $Q$  is a Sylow  $q$ -subgroup of  $A$ . It follows that  $O^r(G) \leq \text{Core}_G(A) = 1$ . In particular,  $G$  is a  $r$ -group and so  $G$  is soluble. Hence,  $A$  is soluble.

We conclude that  $\langle A^G \rangle / \text{Core}_G(A)$  is soluble.

Suppose that  $\mathfrak{S}$  is a Sylow basis of  $G$ . We shall show that  $A^G / \text{Core}_G(A)$  is nilpotent. Without loss of generality we may assume that  $\text{Core}_G(A) = 1$ . Let  $B = \bigcap_q O^q(A)$  be the nilpotent residual of  $A$ . Let  $r$  be a prime dividing  $|G|$  and  $g \in G$ . Then  $g = xy$ , where  $x$  is an element of  $G_r$  and  $y$  is a  $r'$ -element of  $Z = \prod_{q \neq r} G_q$ . It follows that  $B_r = B \cap G_r = O^r(A) \cap G_r$  is a Sylow  $r$ -subgroup of  $B$ . Applying [3, Lemma 1.1.11], we have that  $O^r(A) = O^r(AG_r)$ , which is a normal subgroup of  $AG_r$ , and that  $O^{r'}(A) = O^{r'}(AZ)$ , which is a normal subgroup of  $AZ$ . In particular,  $G_r$  normalises  $O^r(A)$  and  $Z$  normalises  $O^{r'}(A)$ . Moreover,  $B_r$  is contained in  $O^{r'}(A)^y = O^{r'}(A)$ . Since  $G_r$  normalises  $B_r$ , it follows that  $B_r^g = B_r^y$  is contained in  $O^{r'}(A)$  and so it is a subgroup of  $A$ . Consequently, the normal closure  $\langle B_r^G \rangle$  is contained in  $A$  and then  $\langle B_r^G \rangle \leq \text{Core}_G(A) = 1$ . Hence  $B = 1$  and  $A$  is nilpotent.

Therefore,  $\langle A^G \rangle / \text{Core}_G(A)$  is nilpotent.

*Proof (of Theorem 2)* We argue by induction on  $|G|$ . Assume that  $VG_p$  is a proper subgroup of  $G$ . Then  $U \cap VG_p$  is a subnormal subgroup of  $VG_p$ . Since  $UG_p \cap VG_p = (U \cap VG_p)G_p$  is a subgroup of  $G$ ,  $U \cap VG_p$  permutes with the Sylow  $p$ -subgroup  $G_p$  of  $VG_p$ . The induction hypothesis implies that  $G_p$  permutes with  $(U \cap VG_p) \cap V = U \cap V$ . Therefore we may assume that  $G = VG_p$ . An analogous argument with the subnormal subgroup  $U$  of  $UG_p$  and  $V \cap UG_p$  shows that  $G = UG_p$ . Let  $q \neq p$  be a prime dividing  $|G|$  and let  $G_q$  be a Sylow  $q$ -subgroup of  $G$  contained in  $V$ . Then  $G_q \cap U$  is a Sylow  $q$ -subgroup of  $U$  since  $U$  is subnormal in  $G$ . Hence  $G_q$  is contained in  $U$  by order considerations. This means that  $U \cap V$  contains a Sylow  $q$ -subgroup of  $G$  for all primes  $q \neq p$ . Therefore  $|G : U \cap V|$  is a power of  $p$ . Applying [6, Chapter A, Lemma 1.6 (b)], we conclude that  $G = G_p(U \cap V)$  and  $G_p$  permutes with  $U \cap V$ , as required.

*Proof (of Theorem 3)* Suppose that every maximal subgroup of  $G_p \in \mathfrak{S}$  is 3-permutable and that  $G_p$  is not cyclic. By Corollary 7,  $G$  is  $p$ -soluble. Assume that  $G$  is not  $p$ -supersoluble and consider  $G$  of least possible order. Let  $N$  be a minimal normal subgroup of  $G$ . Let  $M/N$  be a maximal subgroup of  $PN/N$ . Then  $M/N = M_1N/N$  for some maximal subgroup  $M_1$  of  $P$ . Since  $M_1$  is 3-permutable, it follows that  $M/N$  is 3 $N/N$  permutable by Lemma 1. Then  $G_pN/N$  has all maximal subgroups 3 $N/N$ -permutable. Assume that  $N$  is a  $p'$ -group. Since  $G_pN/N \cong G_p$ , we have that  $G_pN/N$  is not cyclic and so  $G/N$  is  $p$ -supersoluble by the choice of  $G$ . This implies that  $G$  itself is  $p$ -supersoluble, against the hypothesis. Hence  $N$  is a  $p$ -group. Suppose that  $N$

is contained in  $\Phi(G_p)$ , the Frattini subgroup of  $G_p$ . Then  $G_p/N$  is not cyclic. Hence  $G/N$  is  $p$ -supersoluble by the choice of  $G$ . Moreover,  $N$  is also contained in  $\Phi(G)$ . Since the class of all  $p$ -supersoluble groups is a saturated formation, it follows that  $G$  is  $p$ -supersoluble, contrary to assumption.

Consequently,  $N$  is not contained in  $\Phi(G_p)$ . Let  $M_1$  be a maximal subgroup of  $G_p$  such that  $NM_1 = G_p$ . Let  $q \in \pi(G) \setminus \{p\}$  and let  $G_q$  be the Sylow  $q$ -subgroup of  $G$  in  $\mathfrak{S}$ . Thus  $1 = G_q \cap G_p = G_q \cap NM_1 = (G_q \cap N)(G_q \cap M_1)$ . By [6, Chapter A, Lemma 1.2],  $(N \cap M_1)G_q = NG_q \cap M_1G_q$  is a subgroup of  $G$ . Furthermore,  $(N \cap M_1)G_q \cap N = (N \cap M_1)(G_q \cap N) = N \cap M_1$  is a normal subgroup of  $(N \cap M_1)G_q$ . Hence  $G_q$  normalises  $N \cap M_1$ . On the other hand,  $N \cap M_1$  is a normal subgroup of  $M_1$  and, since  $N$  is abelian, is centralised by  $N$ . Therefore  $N \cap M_1$  is normalised by  $NM_1 = G_p$ . Consequently,  $N \cap M_1$  is a normal subgroup of  $G$  properly contained in  $N$ . Hence  $N \cap M_1 = 1$ . But  $|G_p : M_1| = |NM_1 : M_1| = |N : N \cap M_1| = p$ , hence  $N$  has order  $p$ . If  $M_1$  were not cyclic, then  $G/N$  would be  $p$ -supersoluble by the minimal choice of  $G$ . Thus,  $G$  would be  $p$ -supersoluble, against supposition. Therefore,  $M_1$  is cyclic and  $G/N$  has cyclic Sylow  $p$ -subgroups. This implies that every  $p$ -chief factor of  $G/N$  is cyclic and  $G/N$  is  $p$ -supersoluble. Thus,  $G$  is  $p$ -supersoluble. This final contradiction completes the proof.

*Proof (of Theorem 5)* We prove that  $G$  is  $p$ -soluble by induction on the order of  $G$ . Applying Step 3 of the proof of [14, Theorem 3.3],  $G$  cannot be non-abelian simple. Let  $M$  be a maximal normal subgroup of  $G$ . Assume that  $G_p$  is contained in  $M$ . By Lemma 1,  $\mathfrak{S} \cap M$  is a complete set of Sylow subgroups of  $M$  and every 2-maximal subgroup of  $G_p = M_p \in \mathfrak{S} \cap M$  is  $(\mathfrak{S} \cap M)$ -permutable. By induction,  $M$  is  $p$ -soluble. Furthermore,  $G/M$  is a  $p'$ -group. Thus  $G$  is  $p$ -soluble. Therefore we may assume that  $p$  divides  $|G/M|$ . Then  $M_p = M \cap G_p$  is a proper subgroup of  $G_p$ . Let  $S$  be a maximal subgroup of  $G_p$  containing  $M_p$ . Suppose that  $S$  is cyclic. Then  $G_pM/M$  has a cyclic maximal subgroup. By Lemma 1,  $\mathfrak{S}M/M$  is a complete set of Sylow subgroups of  $G/M$  and every 2-maximal subgroup of  $G_pM/M$  is  $\mathfrak{S}M/M$ -permutable. Therefore, by induction,  $G/M$  is  $p$ -soluble. Furthermore, since  $M_p$  is cyclic, we have that  $M$  is  $p$ -nilpotent by [10, Kapitel IV, Satz 2.8]. Therefore,  $M$  is  $p$ -soluble and so is  $G$ . Hence we may assume that  $S$  is not cyclic. Then  $S$  has two different maximal subgroups which are  $\mathfrak{S}$ -permutable. Thus  $S$  is  $\mathfrak{S}$ -permutable. Let  $q \in \pi(G) \setminus \{p\}$  and let  $G_q$  be the Sylow  $q$ -subgroup of  $G$  in  $\mathfrak{S}$ . It follows that  $M_q = M \cap G_q$  is a Sylow  $q$ -subgroup of  $M$ . Now,  $G_q$  permutes with  $S$  and  $M$ . Applying Theorem 2,  $G_q$  permutes with  $M \cap S = M \cap G_p = M_p$ . Hence,  $M_pM_q = M_p(M \cap G_q) = M \cap M_pG_q$  is a Hall  $\{p, q\}$ -subgroup of  $G$ . By Theorem 9,  $M$  is  $p$ -soluble. Consequently,  $G$  is  $p$ -soluble, as wanted.

*Proof (of Theorem 6)* Suppose that  $G$  is soluble and every 2-maximal subgroup of  $G_p \in \mathfrak{S}$  is  $\mathfrak{S}$ -permutable. Assume, arguing by contradiction, that neither  $G$  is a  $p$ -nilpotent group nor  $G$  has an epimorphic image isomorphic to  $\Sigma_4$ . By Corollary 4 and Theorem 5,  $G$  is  $p$ -soluble.

Let  $N$  be a minimal normal subgroup of  $G$ . The quotient group  $G/N$  inherits the hypothesis of the theorem. Therefore  $G/N$  is  $p$ -nilpotent. Since the class of all  $p$ -nilpotent groups is a saturated formation, it follows that  $N = \text{Soc}(G)$  is a minimal normal subgroup of  $G$  which is complemented in  $G$  by a core-free maximal



$p$ -nilpotent subgroup of  $G$ ,  $M$  say. Moreover,  $C_G(N) = N$  and  $N$  is a  $p$ -group. Hence  $N \leq G_p$ , the Sylow  $p$ -subgroup in  $\mathfrak{3}$ . Then  $G_p = N(G_p \cap M)$  and there exists a maximal subgroup  $M_1$  of  $G_p$  containing  $M_p = G_p \cap M$  such that  $NM_1 = G_p$ . Assume that  $M_p$  is a maximal subgroup of  $G_p$ . Then  $|N| = |G_p : M_p| = p$  and  $G$  is  $p$ -supersoluble. This implies that  $G$  is  $p$ -supersoluble, which contradicts our assumption. Therefore  $M_p$  is not a maximal subgroup of  $G$  and so  $M_p$  is contained in a 2-maximal subgroup  $S$  of  $G_p = NS$ . Let  $p \neq q \in \pi(G)$ . Thus  $1 = G_q \cap G_p = G_q \cap NS = (G_q \cap N)(G_q \cap S)$ . By [6, Chapter A, Lemma 1.2],  $(N \cap S)G_q = NG_q \cap SG_q$  is a subgroup of  $G$ . In particular,  $(N \cap S)G_q \cap N = (N \cap S)(G_q \cap N) = N \cap S$  is a normal subgroup of  $(N \cap S)G_q$  and  $G_q$  normalises  $N \cap S$ . On the other hand,  $N \cap S$  is a normal subgroup of  $G_p$ . Consequently,  $N \cap S$  is a normal subgroup of  $G$  properly contained in  $N$ . Hence  $N \cap S = 1$  and so  $|N| = p^2$ . Since  $C_G(N) = N$ , it follows that  $q$  divides  $p + 1$ . Hence  $p = 2$  and  $G/N$  is isomorphic to  $\Sigma_3$ . Consequently,  $G$  is isomorphic to  $\Sigma_4$ . This contradiction proves the theorem.

Our hypothesis in the next two theorems is that subgroups of  $G_p$  with order  $p$  or 4 (if  $p = 2$ ) are 3-permutable. Let us collect together the arguments common to these two results.

Every subgroup of  $G_p O_{p'}(G)/O_{p'}(G)$  of order  $p$  or 4 (if  $p = 2$ ) is of the form  $TO_{p'}(G)/O_{p'}(G)$  for some subgroup  $T$  of  $G_p$  with order  $p$  or 4 (if  $p = 2$ ). Then, by Lemma 1, every subgroup of  $G_p O_{p'}(G)/O_{p'}(G)$  is  $\mathfrak{3}O_{p'}(G)/O_{p'}(G)$ -permutable. Hence, arguing by induction or minimal counterexample, we assume that  $O_{p'}(G) = 1$ . Hence  $F(G)$ , the Fitting subgroup of  $G$ , is a  $p$ -group.

Assume that  $1 \neq F(G)$  and let  $z$  be an element of  $Z(F(G))$  of order  $p$  and let  $y$  be an element of order  $p$  of  $F(G)$ . Then  $\langle z, y \rangle$  is an elementary abelian subgroup of  $G_p$  and  $G_q$  normalises  $\langle w \rangle$  for each  $w \in \langle z, y \rangle$  and each  $q \neq p$  because  $\langle w \rangle = \langle w \rangle G_q \cap F(G)$  is a normal subgroup of  $\langle w \rangle G_q$ . Hence  $p'$ -elements of  $G$  induce power automorphisms in the abelian socle  $S$  of  $G$ . Applying [3, Lemma 2.1.3], all the  $G$ -chief factors of  $G$  below  $S$  are cyclic and  $G$ -isomorphic.

If  $N$  is a central minimal normal subgroup of  $G$ , then  $\Omega_1(O_p(G))$  is centralised by all  $p'$ -elements of  $G$ . Furthermore, if  $p = 2$ , every subgroup  $Z$  of order 4 of  $F(G)$  is normalised by every  $2'$ -element of  $G$ . Since the automorphism group of  $Z$  is of order 2, it follows that  $O^2(G)$  centralises every subgroup with order 2 and order 4 of  $F(G)$ . In this case, we can apply [10, IV, Satz 5.12], to conclude that  $O^p(G)$  centralises  $F(G)$ .

If  $G$  is  $p$ -soluble, then  $C_G(F(G)) \leq F(G)$  by [10, VI, Hilfssatz 6.5]. Consequently,  $G$  is a  $p$ -group.

*Proof (of Theorem 7)* Assume that all subgroups of  $G_p \in \mathfrak{3}$  with order  $p$  and order 4 (if  $p = 2$ ), with  $G$  a  $p$ -soluble, non- $p$ -supersoluble group of the smallest possible order, are 3-permutable.

By the above arguments,  $p$  is odd,  $O_{p'}(G) = 1$ . Since  $G$  is  $p$ -soluble, it follows that  $S$ , the abelian socle of  $G$ , is just  $\text{Soc}(G)$  and every minimal normal subgroup of  $G$  is not central in  $G$  and has order  $p$ . Let  $N$  be one of them. Then  $C_G(N)$  is a proper normal subgroup of  $G$ . Let  $M$  be a maximal normal subgroup of  $G$  containing  $C_G(N)$ . Since  $N$  has order  $p$ ,  $G/C_G(N)$  is a cyclic group of order dividing  $p - 1$ . In particular,  $|G : M|$  is a  $p'$ -group. Since  $O_{p'}(M) \leq O_{p'}(G) = 1$ , it follows that  $O_{p',p}(M) = O_p(M)$ . The

minimal choice of  $G$  implies that  $M$  is a  $p$ -supersoluble group. Hence  $M/O_p(M)$  is an abelian group of exponent dividing  $p-1$ . Therefore  $O_p(M)$  is a Sylow  $p$ -subgroup of  $G$ . In particular,  $O_p(M) = G_p$  is a normal subgroup of  $G$ .

Since  $G$  is not  $p$ -supersoluble, then  $G$  contains a minimal non- $p$ -supersoluble subgroup  $H$ . Hence  $H$  is one of the groups of [2, Theorem 9]. We will follow the notation of this paper.

Assume that  $|H|$  is divisible only by two primes,  $p$  and  $q$ . Then the Sylow  $p$ -subgroup  $H_p$  of  $H$  is contained in  $G_p$ . The Sylow  $q$ -subgroup  $H_q$  of  $H$  is contained in a conjugate  $G_q^x$  of  $G_q$ , with  $x \in G$ . By taking  $H^{x^{-1}}$  if necessary, we can assume that  $H_q$  is contained in  $G_q$ . Let  $x$  be an element of order  $p$  of  $H$ . Then  $\langle x \rangle G_q$  is a subgroup of  $G$ . Now  $\langle x \rangle G_q \cap G_p = \langle x \rangle (G_q \cap G_p) = \langle x \rangle$  is a normal subgroup of  $\langle x \rangle G_q$ . In particular,  $H_q$  normalises  $\langle x \rangle$ . This rules out the groups of types 2, 4, 6, 8, and 10. Moreover, every element of order a power of  $q$  acts in the same way on all elements of order  $p$ . This rules out the groups of type 3, 5, 7, and 9, since there are elements  $x$  of  $H_p$  such that  $H_q$  does not normalise  $\langle x \rangle$ . Suppose that  $H$  is a group of type 1. If  $s = 1$ , we consider the generator  $c$  of  $C$ , of order  $p$ , which is not normalised by the Sylow  $q$ -subgroup  $H_q$ . If  $s > 1$ , then if  $c$  is a generator of  $C$ ,  $c^{p^{s-1}}$  has order  $p$  and is centralised by  $H_q$ , but  $H_q$  does not centralise the elements of order  $p$  of  $M$ . This contradicts the fact that  $H_q$  induces the same automorphism on all cyclic subgroups of order  $p$ .

Assume now that  $H$  has order divisible by three primes,  $p$ ,  $q$ , and  $r$ . Then  $H$  is one of the groups of types 11 or 12. As above, we can assume that the Sylow  $q$ -subgroup  $H_q$  is contained in  $G_q$ . As before,  $G_q$  normalises  $\langle x \rangle$  for each  $x \in G_p$ , in particular,  $H_q$  normalises  $\langle x \rangle$  for each  $x \in H_p$ . If  $G$  is a group of type 12, then  $H_p = P$  is an extraspecial group of order  $p^3$  and exponent  $p$  and the elements of  $H_q = M$  act on the cyclic subgroups of  $P$  in the same way. This is impossible since  $M$  does not centralise  $P$ . Assume that  $G$  is a group of type 11. There exists  $z \in G$  such that the Sylow  $r$ -subgroup  $H_r$  of  $H$  is contained in  $G_r^z$ . Let  $c$  be the generator of  $C$ . Given  $y \in G_r$ , there exists an integer  $t(y)$  such that if  $x$  is an element of order  $p$  of  $G_p$ ,  $x^y = x^{t(y)}$  for each  $y \in G_r$ . In particular, given an element  $x$  of  $H_p^z$ ,  $x^c = x^{t(c)}$ . Since  $c$  acts in the same way on all elements of  $G_p$ , for every element  $x$  of  $H_p$ ,  $x^c = x^{t(c)}$ . But this implies that  $MC$  acts as a group of power automorphisms on  $P$ , in particular,  $MC$  acts as an abelian group on  $P$ . This implies that  $H$  is  $p$ -supersoluble, a contradiction.

*Proof (of Theorem 8)* Let  $G$  be a group in which every cyclic subgroup with order  $p$  or order 4 (if  $p = 2$ ) of  $G_p$  is  $\mathfrak{Z}$ -permutable. Assume that the order of  $G_p$  is greater than  $p$ . We prove that  $G$  is  $p$ -soluble by induction on the order of  $G$ . Applying the above arguments, we may assume that  $O_{p'}(G) = 1$  and every abelian minimal normal subgroup of  $G$  is of order  $p$ .

Let  $M$  be a maximal normal subgroup of  $G$ . Then, by Lemma 1,  $M$  satisfies the hypotheses of the theorem. Therefore either  $M_p$  is of order  $p$  or  $M$  is  $p$ -soluble. If  $M$  is  $p$ -soluble, then  $M$  is  $p$ -supersoluble by Theorem 7. Since  $O_{p'}(M) \leq O_{p'}(G) = 1$ , it follows that  $M_p = G_p \cap M$  is a normal Sylow  $p$ -subgroup of  $M$  by [3, Lemma 2.1.6].

Let  $A$  be a maximal normal subgroup of  $G$  such that  $A \neq M$ . Then  $G = AM$ . Applying [3, Theorem 1.1.19], there exist Sylow  $p$ -subgroups  $A_p$  and  $M_p$  of  $A$  and  $M$ , respectively, such that  $A_p M_p$  is a Sylow  $p$ -subgroup of  $G$ . If  $|A_p| = |M_p| = p$ , then

$|G_p| = p^2$  and, by Theorem 3,  $G$  is  $p$ -supersoluble. Suppose that the order of  $M_p$  is greater than  $p$ . Then  $M_p$  is normal in  $G$ . If  $A$  were  $p$ -supersoluble, then  $A_p$  would be also normal in  $G$  and so would be  $G_p$ . Then  $G$  is  $p$ -soluble. Assume that  $|A_p| = p$ . Then  $M_p$  is not contained in  $A$  and so  $G = AM_p$ . This means that  $G/A$  is of order  $p$  and  $|G_p| = p^2$ . By Theorem 3,  $G$  is  $p$ -supersoluble. Therefore  $G$  is  $p$ -soluble.

Therefore we may assume that  $M$  is the unique maximal normal subgroup of  $G$ . Assume that  $MG_p$  is a proper subgroup of  $G$ . Then  $\mathfrak{Z} \cap MG_p$  is a complete set of Sylow subgroups of  $MG_p$  and every subgroup of  $G_p$  with order  $p$  or order 4 (if  $p = 2$ ) is  $(\mathfrak{Z} \cap MG_p)$ -permutable. By induction,  $MG_p$  is  $p$ -soluble and  $M_p$  is a normal subgroup of  $G$ . Since  $O_{p'}(G) = 1$ , it follows that  $M_p \neq 1$  and  $F(G) = F(M)$  is a non-trivial  $p$ -group. Suppose that  $p = 2$ . Since  $M$  is 2-soluble, we conclude that  $M$  is a 2-group and so  $G$  is 2-soluble. Assume that  $p$  is odd and every abelian minimal normal subgroup of  $G$  is non-central. Let  $N$  be one of them. Then  $C_G(N)$  is contained in  $M$  and so  $G/M$  is a  $p'$ -group. Consequently,  $M_p$  is a normal Sylow  $p$ -subgroup of  $G$  and  $G$  is  $p$ -soluble.

We may therefore assume that  $G = MG_p$  and  $|G : M| = p$ . If  $M$  were not  $p$ -soluble, then  $M_p$  must be of order  $p$  and so the order of the Sylow  $p$ -subgroups of  $G$  would be  $p^2$ . By Theorem 3,  $G$  is  $p$ -supersoluble.

Consequently, in all cases,  $G$  is  $p$ -soluble and the induction argument is complete.

*Proof (of Corollary 5)* Assume that  $G$  is a group in which every cyclic subgroup with order  $p$  or order 4 (if  $p = 2$ ) of  $G_p$  is 3-permutable. If the order of  $G_p$  is greater than  $p$ , then  $G$  is  $p$ -soluble by Theorem 8. If the order of  $G_p$  is  $p$ , then  $G$  has Hall  $\{p, q\}$ -subgroups for all  $q \in \pi(G)$ . By Theorem 9,  $G$  is  $p$ -soluble. In both cases, we have that  $G$  is  $p$ -soluble. Applying Theorem 7,  $G$  is  $p$ -supersoluble.

## Acknowledgements

This Project was funded by the Deanship of Scientific Research (DSR), at King Abdulaziz University, Jeddah, under grant no. (1/31/RG). The authors, therefore, acknowledge with thanks DSR technical and financial support. The second and the third authors have been supported by the research project MTM2010-19938-C03-01 from the *Ministerio de Ciencia e Innovación*, Spain, and the first author has also been supported by the National Natural Science Foundation of China (Grant No. 11271085).

We also thank E. Vdovin for providing us with a proof of Theorem 9.

## References

1. Asaad, M., Heliel, A.A.: On permutable subgroups of finite groups. *Arch. Math. (Basel)* **80**, 113–118 (2003). DOI 10.1007/s00013-003-0782-4
2. Ballester-Bolínches, A., Esteban-Romero, R.: On minimal non-supersoluble groups. *Rev. Mat. Iberoamericana* **23**(1), 127–142 (2007). DOI 10.4171/RMI/488
3. Ballester-Bolínches, A., Esteban-Romero, R., Asaad, M.: Products of finite groups, *de Gruyter Expositions in Mathematics*, vol. 53. Walter de Gruyter, Berlin (2010). DOI 10.1515/9783110220612. URL <http://dx.doi.org/10.1515/9783110220612>
4. Deskins, W.E.: On quasinormal subgroups of finite groups. *Math. Z.* **82**, 125–132 (1963). DOI 10.1007/BF01111801

5. Doerk, K.: Eine Bemerkung über das Reduzieren von Hallgruppen in endlichen auflösbaren Gruppen. *Arch. Math. (Basel)* **60**, 505–507 (1993). DOI 10.1007/BF01236072
6. Doerk, K., Hawkes, T.: *Finite Soluble Groups, De Gruyter Expositions in Mathematics*, vol. 4. Walter de Gruyter, Berlin, New York (1992). DOI 10.1515/9783110870138. URL <http://dx.doi.org/10.1515/9783110870138>
7. Hall, P.: On the Sylow systems of a soluble group. *Proc. London Math. Soc. (2)* **43**, 316–323 (1937). DOI 10.1112/plms/s2-43.4.316
8. Heliel, A.A., Al-Gafri, T.M.: On conjugate- $\mathfrak{J}$ -permutable subgroups of finite groups. *J. Algebra Appl.* **12**(8), 1350.060 (14 pages) (2013). DOI 10.1142/S0219498813500606
9. Heliel, A.A., Li, X., Li, Y.: On  $\mathfrak{J}$ -permutability of minimal subgroups of finite groups. *Arch. Math. (Basel)* **83**, 9–16 (2004). DOI 10.1007/s00013-004-1014-2
10. Huppert, B.: Endliche Gruppen I, *Grund. Math. Wiss.*, vol. 134. Springer Verlag, Berlin, Heidelberg, New York (1967)
11. Kegel, O.H.: Sylow-Gruppen und Subnormalteiler endlicher Gruppen. *Math. Z.* **78**, 205–221 (1962). DOI 10.1007/BF01195169
12. Li, X., Li, Y., Wang, L.:  $\mathfrak{J}$ -permutable subgroups and  $p$ -nilpotency of finite groups II. *Israel J. Math.* **164**, 75–85 (2008). DOI 10.1007/s11856-008-0021-6
13. Li, Y., Heliel, A.A.: On permutable subgroups of finite groups II. *Comm. Algebra* **33**(9), 3353–3358 (2005). DOI 10.1081/AGB-200058541
14. Li, Y., Li, X.:  $\mathfrak{J}$ -permutable subgroups and  $p$ -nilpotence of finite groups. *J. Pure Appl. Algebra* **202**, 72–81 (2005). DOI 10.1016/j.jpaa.2005.01.007
15. Li, Y., Wang, L., Wang, Y.: Finite groups with some  $\mathfrak{J}$ -permutable subgroups. *Glasgow Math. J.* **52**, 145–150 (2010). DOI 10.1017/S0017089509990231
16. Vdovin, E.P., Revin, D.O.: Theorems of Sylow type. *Russian Math. Surveys* **66**(5), 829–870 (2011). DOI 10.1070/RM2011v066n05ABEH004762
17. Wang, L.F., Wang, Y.M.: A remark on  $\mathfrak{J}$ -permutability of finite groups. *Acta Math. Sinica* **23**(11), 1985–1990 (2007). DOI 10.1007/s10114-005-0906-9