

Mental fatigue impairs repeated Sprint and jump performance in team sport athletes

Walter Staiano, Luis Raimon Salazar Bonet, Marco Romagnoli, Christopher Ring



PII: S1440-2440(23)00458-9

DOI: <https://doi.org/10.1016/j.jsams.2023.10.016>

Reference: JSAMS 2890

To appear in: *Journal of Science and Medicine in Sport*

Received date: 4 March 2023

Revised date: 19 October 2023

Accepted date: 22 October 2023

Please cite this article as: W. Staiano, L.R.S. Bonet, M. Romagnoli, et al., Mental fatigue impairs repeated Sprint and jump performance in team sport athletes, *Journal of Science and Medicine in Sport* (2023), <https://doi.org/10.1016/j.jsams.2023.10.016>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Mental Fatigue Impairs Repeated Sprint and Jump Performance in Team Sport Athletes

Submission Type: Original Investigation

Authors: Walter Staiano^{1,2}, Luis Raimon Salazar Bonet³, Marco Romagnoli¹, Christopher Ring⁴

¹ Department of Physical Education and Sport, University of Valencia (SP)

² Department of Psychology, Biological and Cognitive Psychology, University of Southern Denmark (DK)

³ International University SEK, (EC)

⁴ School of Sport, Exercise & Rehabilitation Sciences, University of Birmingham (UK)

Corresponding Author :

Dr. Walter Staiano

Department of Physical Education and Sport, University of Valencia, Valencia, Spain.

Address: C/ Gascó Oliag, 3, 46010, Valencia (Spain).

Tel: +34663262239

Email: walterstaiano@gmail.com

Preferred Running Head: Mental fatigue and sprint and jump performance

Abstract Count: 250

Text-only word count: 4059

Figures and Tables: 1 Table and 3 Figures

Mental Fatigue Impairs Repeated Sprint and Jump Performance in Team Sport Athletes

Submission Type: Original Investigation

Preferred Running Head: Mental fatigue and repeated sprint and jump performance

Abstract Count: 250

Text-only word count: 4355

Figures and Tables: 1 Table and 3 Figures

Abstract

Objectives: We tested whether mental fatigue (MF), induced by a cognitively-demanding task, would impair repeated sprint ability (RSA) and repeated jump ability (RJA) performance, and whether physical fatigue and MF would impair psychomotor vigilance. **Design:** Randomized within-participant design. **Methods:** After establishing baseline peak countermovement jump (CMJ), 18 male participants performed 12 maximal 20-m (10-m linear + 10-m directional) repeated sprints (RSA random test) followed by 12 maximal repeated CMJs (RJA test) subsequent to 30-min Stroop task (MF) or a documentary (Control). Peak and mean running time and height, percent decrement score (S_{dec}), blood lactate, heart rate and RPE were measured for CMJ, RSA, and RJA tests. MF (M-VAS) and psychomotor vigilance [psychomotor vigilance test (PVT)] were measured at baseline, after each condition, and after the RSA/RJA tests. **Results:** Compared to Control, the Stroop task elevated MF ($p = .001$), RPE ratings (all $p < .031$), and mean and S_{dec} performance in directional (but not linear) RSA (all $p < .032$) and RJA tests (all $p < .034$). PVT score worsened after Stroop task ($p = .011$) but not Control, declined after RSA/RJA tests in both conditions (all $p < .023$) and was lower in the MF condition ($p = .029$). No condition differences were noted for peak (CMJ, RSA and RJA tests) performance, blood lactate, and heart rate. **Conclusion:** MF impairs directional RSA, and RJA performance. This impairment was linked with increased RPE and without physiological changes. The progressive impairment in PVT score suggests a cumulatively negative effect of mental and physical fatigue on psychomotor vigilance.

Keywords: anaerobic exercise, cognitive load, cognitive performance, sport performance, physical fatigue, strength endurance

INTRODUCTION

Mental fatigue (MF) is a psychobiological state that can be induced by cognitively demanding tasks, characterized by subjective tiredness or lack of energy and impaired cognitive performance¹. Sport scientists have confirmed that MF can impair aerobic endurance performance^{1,2} as well as sports-specific psychomotor performance, defined as highly complex motor behavior that results from the cognitive processing of sensory and perceptual information in a sport-specific context³.

On the other hand, evidence on the effect of MF on anaerobic performance and strength is less explored. Reviews conclude that MF does not affect physical activities that require brief bursts of strength, such as sprints, jumps, and maximal contractions^{1,3}. Indeed, studies report that countermovement jumps (CMJ) and running sprints are unaffected by MF^{4,5}. Other studies note that MF does not impair short-lasting (< 3 min) strength/power and sprint exercises^{4,6}. For instance, linear repeated sprint ability (RSA) performance was unaffected by MF⁷. However, another line of investigation concerning strength endurance proved otherwise, with studies showing that MF can impair submaximal strength endurance exercise^{8,9,10}. For instance, when performing submaximal contractions repeatedly at 25% of maximum voluntary contraction, MF increased the time to exhaustion⁸. Other research, including a recent review on strength endurance, showed that MF reduced the number of repetitions and sets of strength exercise performed^{9,10}. It can be argued that strength endurance, similarly to aerobic endurance performance, is susceptible to MF due to higher cognitive components of pacing, decision-making, and, perceived effort moderation^{1,3}. In addition, it is worth considering that such cognitive components are more relevant when the physical demand is accompanied by the situational demands that players experience when processing information before deciding where and how to move, while also considering their physical demands, such as postural control and balance stabilization¹². In this complex environment, the negative impact of MF could be exacerbated, with greater impairments of decision making, ball passing, dribbling accuracy, directional changes, reactive balance and stability^{1,3,11,12}.

Therefore, it is worth noting that evidence to date on the effect of MF on anaerobic performance and strength endurance are restricted to studies that examined the effects of MF on either isolated anaerobic physical tests, such as CMJ, or on physical tests that do not incorporate complex cognitive components, such as decision-making, pacing, directional changes, and balance control, which are important features of sport performance. Accordingly, studies of the effects of MF on anaerobic strength and strength endurance could/should consider how multiple elements of complex situations (e.g., situation demands and personal capabilities) might interact to determine performance outcome in sport¹³. In other words, to increase ecological validity, protocols could examine the effects of mental and physical fatigue on subsequent repeated performance of tasks that require control of posture/balance as well as decision making and visuo-motor reactions. Scarce examples of this approach can be found in the literature^{1,3}. First, the only study using a repeated jump ability (RJA) test explored whether RJA is affected by the environmental factor heat stress¹⁴. Secondly, performance of a repeated sprint ability (RSA) test was only tested linearly and, notably, no study has explored the effect of MF on the RSA test with a linear and directional component combined. Previously, this specific RSA test with both of these components, was used in a training study and was sensitive to mental fatigue-inoculation cognitive training¹⁵. In addition, in sports, such as football,

basketball and volleyball, athletic actions, such as high intensity sprinting, changing direction, jumping, accelerating, and decelerating (which are typically examined only from a physiological perspective), have a negative impact on cognition and in particular in vigilance and sustained attention¹⁶. Despite such impact, there is scarce evidence of the impact of physical fatigue and the combination of physical and mental fatigue on cognitive performance.

Therefore our study was designed to address these gaps in the literature and its purposes were twofold. Our first purpose was to determine the effect of a highly demanding cognitive task inducing MF on subsequent repeated sprinting and jumping performance using a protocol with multiple elements to increase complexity, in terms of cognitive demands (e.g., sprinting with changes of direction) and exercise demands (e.g., completing repetitive bouts with incomplete recovery). We hypothesized that a 30-min response inhibition cognitive task would induce MF and impair peak and mean time of the directional component but not the linear component of RSA. Furthermore, we hypothesized that MF would not affect peak height during isolated CMJs (measured at baseline and after each experimental treatment) and the RJA, whereas MF would impair mean height and fatigue index (measured as decrement score) during the RJA. Our second study purpose was to determine the serial effects of MF alone and combined with physical fatigue on the performance of a psychomotor vigilance test (PVT). Previous studies have shown that high intensity aerobic endurance exercise can impair subsequent cognitive performance^{7,16}. Together with evidence that MF impairs cognitive performance⁷, this literature highlights a shared neurocognitive connection concerning the regulation of mental and physical effort. Here we hypothesized that reaction times during the PVT would be slower following 30-min of a mentally fatiguing task compared to watching neutral videos, and slowest following the serial combination of MF plus physical fatigue.

METHODS

Eighteen healthy and fit male [mean $\pm$ SD, age 25 ± 5 years, height 181 ± 5 cm, weight 75 ± 8 kg, peak oxygen uptake 55 ± 4 ml $\cdot$ kg⁻¹ $\cdot$ min⁻¹] participants were recruited from amateur basketball (n = 6), handball (n = 7) and football (n = 5) clubs. They had at least three years playing experience and were classified as performance level 2 (trained/development)¹⁷. Given that we only had access to male clubs, we were not able to test female participants. They signed an informed consent and the study protocol was approved by the Ethics Committee of the School of Sport and Physical Education at Valencia University in accordance with the Declaration of Helsinki. All participants were unaware of the study purposes and hypotheses.

This study employed a randomized counterbalanced within-participant experimental design. Participants completed 3 laboratory sessions: a familiarization session followed by two counterbalanced (mental fatigue, control) testing sessions (Fig. 1). Time of testing, environmental conditions, and exercise equipment settings were standardized. At least 10 days separated testing sessions to allow recovery from potential muscle damage. Participants were instructed to sleep for at least 7 h, drink 35 ml of water per kilogram of body weight, refrain from alcohol and caffeine, and avoid any vigorous exercise 24 h prior each laboratory session. Their compliance with instructions was checked and verified. Participants declared any

medication/drug use and acute illness, injury, or infection. Time of testing, environmental conditions, meal times and equipment settings were standardized.

During the familiarization session, participants were provided with standardized instructions for anchoring of Borg 6-20 rating of perceived exertion (RPE) and completed the RSA random test¹⁸, counter movement jump test¹⁴, and RJA test¹⁴. Furthermore, they were familiarized with the Stroop task, PVT¹⁹, and the psychological and physiological measures. During the testing sessions, participants rated their motivation to complete the whole testing session, and MF, warmed-up, and performed the baseline CMJ test, which comprised three vertical jumps on a force plate (Kistler, Arizona, US) separated by 20 s rest. Afterwards they performed a 10-min PVT using a computer running E-prime 2.0 (Pittsburg, USA) software to assess sustained attention and vigilance to perform. Next, they completed one of two 30-min experimental conditions while sitting comfortably in a quiet, dimly lit room. In the MF condition they performed an incongruent Stroop colour-word task using a tablet running the SOMA-NPT app (Sswitch, Lucerne, Switzerland). In the control condition (CON) they watched an emotional neutral documentary about cars on the same screen¹. At the end of each condition, they completed the National Aeronautics and Space Administration Task Load Index (NASA-TLX)²⁰, performed a 10-min PVT, and rated MF. After warming-up, they completed another CMJ test, rested 5-min, provided a blood sample, rated their motivation to complete the whole testing session, warmed-up for 10-min, and completed the RSA random test. This running test measures linear acceleration, change of direction, visuomotor response, and decision making. The test comprised of 12 sprints, with each sprint followed by 10 s active recovery while jogging 20 m to the starting position. Each sprint comprised a 10-m linear sprint plus a 10-m directional sprint to 1 of 3 randomly cued locations (straight or 45 degrees to the right or to the left). The location of each directional sprint was cued by the illumination of 1 of 3 colored lights after completing the previous 10-m linear sprint. We used a FitLight Trainer system (Fitlight Corp, Ontario, Canada) to measure linear and directional sprints and to cue directions. Performance was measured as peak and mean time taken to complete the 10-m linear acceleration sprint (time [s]) and 10-m directional sprint (time [s]). HR and RPE were measured after every sprint. At completion of the test, participants rested for 10-min and provided a blood sample. They then warmed-up for 5-min and completed the RJA test. This jumping test comprised 12 maximal jumps on a force plate (Kistler, Arizona, US) separated by 5 second intervals. Participants were instructed to jump from a stationary standing position, to use a preparatory countermovement consisting of a 90° knee flexion, and to keep their hands on their waist¹⁴. Finally, they completed a NASA-TLX and performed a 10-min PVT. All warm-up protocols preceding the CMJ, RSA and RJA tests were standardized across experimental visits.

For the CMJ test, the peak force and height for each of the three jumps were recorded, and the two closest values were averaged. For the RSA random test, peak performance time, mean performance time, and percentage decrement score (S_{dec}) were measured. For the RJA test, peak height, mean height, and S_{dec} were computed. S_{dec} is a measure of fatigue expressed as percentage of decline in maximal performance and it is computed using equation 2 for RJA and equation 3 for RSA as formulated by Girard and colleagues.²¹ For the PVT, mean reaction time (ms) for responses between 100 and 500 ms, and number of lapses, defined as responses slower than 500 ms, were computed¹⁹.

Heart rate was measured using a telemetric chest sensor (Polar H10, Polar Electro Oy, Kempele, Finland) during the RSA random test, specifically at the end of each sprint completed. Blood lactate concentration (mmol/l) was measured by taking a 5- μ l sample of whole fresh capillary blood from the right middle finger and analyzed using a portable analyzer (Lactate Pro LT-1710, Arkray, Shiga, Japan). Samples were collected at baseline before each

condition, before the RSA and before RJA tests. A 6-20 Borg scale²² was used to assess perceived effort during each sprint of the RSA random test, and overall effort across the RJA test.

Motivation to complete the testing session was measured by rating the statement “I am motivated to perform the test” using a 5-point Likert scale, with anchors of 0 (not at all) and 4 (extremely). Subjective workload after each condition and physical test was measured using the mental, physical demand, frustration and effort subscales of the NASA-TLX²⁰. MF before and after each condition was measured using a specific visual analogue scale (M-VAS): participants indicated how mentally tired they were on a scale, anchored by “not all exhausted” and “completely exhausted”, by placing a mark on a 10-cm line.

All data are presented as mean $\pm$ SD. After testing for normality using Shapiro–Wilk test, histograms, Q–Q plots, and box-plots, RSA and RJA performance parameters, blood lactate measures, motivation data, NASA TLX, and RPE after the RJA test were analyzed using paired sample t tests. M-VAS scales and CMJ parameters were analyzed using 2 condition (MF, Control) by 2 time (pre 30-min manipulation, post 30-min manipulation) ANOVAs. HR and RPE (during the RSA) were analyzed using 2 condition (MF, Control) by 12 sprints (1 ... 12) ANOVAs. PVT reaction time and lapses were analyzed using 2 condition (MF, Control) by 3 time (pre 30-min manipulations, pre RSA test, post RJA test) ANOVAs. Interaction effects were followed up with relevant corrected pairwise comparisons using the Bonferroni method (post hoc analysis) for simple main effects, otherwise main effects are reported. Significance was set at 0.05 (2-tailed) for all analyses. The effect sizes for the ANOVAs were calculated as partial eta squared (η^2p), with the small = .02, medium = .13 and large = .26 interpretation. For t tests, Cohen’s d was calculated as effect size, with the small = .20, medium = .50 and large = .80 interpretation. Data analysis was conducted using the Statistical Package for Social Sciences (SPSS 27).

RESULTS

In the RSA random test (Fig. 2A), mean running time was significantly longer for MF than control during the directional phase ($p = .032$, $d = 0.491$) but not the acceleration phase ($p = .192$, $d = 0.112$). Moreover, running S_{dec} was significantly greater ($p = .027$, $d = 0.511$) in the MF condition compared to control (Fig. 2C). Similarly, in the RJA test (Fig. 2B), the MF condition was characterized by significantly poorer jumping, namely shorter mean height ($p = .009$, $d = 0.599$), and greater S_{dec} ($p = .034$, $d = 0.523$) (Fig. 2B), compared to the control condition.

In contrast, in the RSA random test, no significant condition differences were detected for peak running time during the acceleration ($p = .371$, $d = 0.291$) and directional ($p = .271$, $d = 0.211$) phases (Fig. 2A). In the case of jumping performance, there were no significant condition ($p = .581$, $\eta^2p = .10$; $p = .272$, $\eta^2p = .11$), time ($p = .183$, $\eta^2p = .10$; $p = .212$, $\eta^2p = .11$), or condition by time ($p = .321$, $\eta^2p = .10$; $p = .441$, $\eta^2p = .06$) effects for peak height during the CMJ test (Fig. 2B). Moreover, we failed to detect any significant condition differences in peak height ($p = .217$, $d = 0.141$) during the RJA test (Fig. 2B).

The impact of MF on vigilance was assessed by performance on the PVT. Two condition by three time ANOVAs generated significant interaction effects for RT ($p = .021$, $\eta^2p = .18$). Follow-up tests revealed no significant condition differences at baseline ($p = .531$) whereas reaction time slowed significantly relative to baseline after the MF manipulation only ($p = .011$), which was slower ($p = .002$) than CON. Moreover, after the physical tests, reaction time

significantly slowed in both MF ($p = .015$) and CON ($p = .023$) conditions, and was slower in the MF ($p = .029$) compared to control condition (Fig. 2D). No significant interaction or main effects were noted for lapses.

Regarding heart rate, ANOVA yielded a significant main effect of time, confirming that heart rate ($p < .001$, $\eta^2p = .91$) increased over the 12 sprints. Importantly, no significant condition ($p = .291$, $\eta^2p = .10$) or condition by time ($p = .237$, $\eta^2p = .09$) effects were found (Fig. 3A). Similarly, no significant condition differences in blood lactate were observed before the manipulations ($p = .321$, $d = 0.181$), before the RSA random test ($p = .211$, $d = 0.154$) and before the RJA test ($p = .611$, $d = 0.222$) (Fig. 3B).

RPE during the RSA random test yielded a significant condition by time interaction ($p = .031$, $\eta^2p = .18$); effort increased over the 12 sprints in both conditions ($p < .001$, $\eta^2p = .44$), and running was more effortful during the MF condition ($p = .022$, Fig. 3C). Similarly, effort at completion of the RJA test was significantly higher in the MF condition compared to CON ($p = .017$, $d = 0.698$) (Fig. 3D).

Motivation did not significantly differ between conditions before the manipulations ($p = .422$, $d = 0.279$) and physical tests ($p = .515$, $d = 0.176$). The NASA TLX subscales completed after the manipulations revealed significantly higher mental demand ($p < .001$, $d = 0.999$) and effort ($p = .031$, $d = 0.713$) during the MF than CON condition. No significant condition difference was noted for physical demand ($p = .519$, $d = 0.154$) and frustration ($p = .714$, $d = 0.231$) (Table 1). The NASA TLX subscales completed after the physical tests revealed significantly higher mental demand ($p = .021$, $d = 0.375$), effort ($p = .035$, $d = 0.272$), and frustration ($p = .037$, $d = 0.455$) during the MF than CON condition. No significant condition difference was noted for physical demand ($p = .813$, $d = 0.199$) (Table 1). A 2 condition by 2 time ANOVA on the M-VAS ratings produced an significant interaction effect ($p = .001$, $\eta^2p = .31$); MF increased from pre (1.3 ± 0.5) to post (7.2 ± 1.1) cognitive task ($p = .021$), whereas MF did not change from pre (1.0 ± 0.7) to post (2.4 ± 0.8) video task ($p = .713$).

DISCUSSION

Our study examined whether a state of MF, induced by a highly demanding cognitive task, would impair repeated sprinting (RSA) and jumping (RJA) performance, increase rating of perceived exertion (RPE), and impair sustained attention and vigilance (PVT). In line with our hypotheses, we found that MF impaired mean running time and running S_{dec} during the directional phase of the RSA test as well as mean jumping height and S_{dec} during the RJA test. However, no differences were detected for peak running time (RSA random test) and height (CMJ and RJA tests). Moreover, PVT performance was affected by MF and combined physical and mental fatigue. These key findings are discussed below.

Our first study purpose was to determine the effects of MF on subsequent repeated sprinting and jumping performance. In partial support of our hypothesis that MF would impair repeated exercise performance, we found that a state of MF, induced by performing a highly demanding 30-min classic Stroop response inhibition task, slowed mean running time by 9% during the directional phase of the RSA random test compared to control. This finding is in line with another study showing that fatigue-inoculation training improved intermittent running performance during the directional phase of the RSA random test¹⁵. Furthermore, it

corroborates previous studies showing that mental fatigue is associated with poorer physical and technical performance in football and decreased decision-making skill and visual search performance in basketball^{1,3,11}. Taken together, these findings provide evidence that the complex cognitive requirement to respond to a visual stimulus and decide which direction to take while sprinting is slowed when mentally fatigued. We also found that MF increased the S_{dec} during the RSA random test, a measure of sprinting fatigue over time. This evidence demonstrates that MF also reduced resilience to physical fatigue during repeated high intensity running, and it is in line with previous studies on the negative effects of MF on shuttle and yo-yo tests. The present data corroborate the argument that Yo-Yo Tests and intermittent high-intensity running tests in soccer players and other team sport athletes are affected by mental fatigue by altering their perception of effort²³. As hypothesised, we failed to observe any effect of MF on peak time, mean time, and S_{dec} during the linear acceleration phase of the RSA random test. These null findings for basic high intensity running (i.e., sprinting) in a straight line, add to those yielded by a previous study that also failed to detect any effect of MF on repeated straight line sprinting in footballers^{1,7}. These null findings argue that high intensity linear sprinting performance, which in general relies on energy supply, metabolic accumulation, reduced neural drive and environmental factors, is immune to the deleterious effects of MF^{1,7}.

Similarly, and in partial support of our hypothesis, we found that MF impaired performance of the 12-jump RJA test, whereas MF did not impair performance of the 3-jump CMJ test. Jump height during the CMJ tests did not differ between the MF and control conditions. This null finding is broadly in agreement with previous studies^{4,5} that reported no effect of MF on single CMJ performance. Importantly, when participants were required to execute a series of jumps, with incomplete recovery between jumps, as required by our RJA protocol, we now found that MF impaired jumping mean height by 8% and S_{dec} by 58% compared to control. Our RJA results are broadly compatible with those showing that mean (but not peak) height during a series of 5 repeated jumps was impaired by heat stress¹⁴. Our RJA protocol comprising 12 jumps, proved that repeated jumping is sensitive to increased cognitive demands faced by the performer. Although we did not perform any biomechanical analysis, we could speculate that MF impacted the control of posture². It is possible that MF impaired the ability of participants to balance based on external environmental cues. In this case, the taking off and landing phases of jumping may be impaired if insufficient recovery is granted to allow proper readjustment. MF may have impaired the ability to create faster adjustment of reactive balance and adaptability²⁴. This explanation could account for why peak height was not affected by MF whereas mean height and S_{dec} were impaired by MF. It is worth noting that the impaired S_{dec} due to MF in the present study is in line with previous studies showing that MF affected the number of repetitions and volume-load ratio in resistance exercise and training to exhaustion^{9,10,25}. Although our protocols were submaximal and not until exhaustion, it is evident from the S_{dec} that MF would have affected the number of repetitions if exercise had carried on to exhaustion. Overall, the present study shows that MF affected the higher order components required to perform the RSA and RJA tests, namely, the directional phase of repeated running and the repeated jumps with uncomplete recovery.

Our second study purpose was to determine the effects of mental and physical fatigue on sustained attention and vigilance as measured by the psychomotor vigilance test. In support of our hypothesis, we found that reaction times were 18% slower following the 30-min Stroop task compared to control. These results, indicating that a state of MF reduced vigilance task performance, are in line with previous studies assessing the effect of MF on cognitive performance^{7,19}. This behavioural measure of MF provided by the PVT is corroborated by the

subjective MF and NASA TLX ratings, which confirmed that the Stroop task was more mentally fatiguing, mentally demanding, and effortful than watching emotionally neutral videos. Also, in line with our hypothesis, we found that reaction times were slowest after completion of the exercise tasks. Previous studies show that exercise can have positive, null, or negative effects on cognition depending on the exercise dose²⁶. In line with previous research¹⁶, here we found that high intensity exercise impaired reaction times during a vigilance task. That the MF condition slowed reaction time during the PVT more than the control condition after the RSA and RJA tests establishes a cumulative effect of mental and physical fatigue on participants' sustained attention and vigilance to perform. Indeed, MF plus physical fatigue slowed reaction time 16% more compared to physical fatigue alone. These results are in line with the only two previous studies, to our knowledge^{7, 27}, which illustrate the additive negative effects of MF and physical fatigue on reaction times during a vigilance task. Vigilance is an important cognitive element in sport performance, and, therefore, the present findings highlight the importance of considering the cumulative effects of mental and physical fatigue on athletic performance.

It is worth mentioning that HR and blood lactate did not differ between conditions. Accordingly, these metabolic and cardiac factors are unlikely to mediate the decline in performance of the RSA and RJA tests observed in the MF condition. These results are in line with studies of MF and sport performance, and underscore the view that physiological measures are not affected by MF and they do not change during the RSA and RJA performance^{1,3}. Although we did not take any biomechanical measure, we speculate that MF may impact motor control and intra/inter-muscular coordination of complex movement, alter volitional, reactive balance and postural control and impair one's ability to maintain and regain stability¹². The RPE measured during the RSA random test and at completion of the RJA test confirmed the increased effort associated with performing the physical tests and are in line with the previous literature indicating that RPE may be a main mediator between MF and sport performance^{1,3}. The RPE scale in this paradigm represented the effort needed to sprint and jump repetitively. Participants in the MF condition rated their effort to complete the RSA and RJA tests to be between 4% and 8% more effortful than control, which supported the hypothesis that MF increased RPE. The available data demonstrate that MF negatively impacts the perceived effort to perform. It appears in our study as in previous ones⁴ that MF has no effect on peak time and height. However, when running and jumping exercises are performed repetitively and with more complex cognitive demands, it makes these exercises more effortful.

The present study helps clarify the relationship between MF and repeated anaerobic performance/strength endurance, an important component in team sport²³. However, studies using strength endurance components and MF, biomechanical measures, and a range of athletic levels are necessary to better understand the effect of MF on postural control, and to clarify how the level of the athletes impacts resilience toward mental fatigue. Indeed, a limitation of the present study was the lack of in depth measures and analysis of biomechanical postural control which would have better elucidated the biomechanical postural mechanisms affected by MF.

Future studies should evaluate short-term recovery strategies to reduce MF such as binaural sounds²⁸ or psychological and nutritional interventions²⁹. Alternatively, coaches could train athletes to cope better with MF using brain endurance training^{15,30}, a form of fatigue-inoculation training.

CONCLUSION

Our study showed that a state of MF, induced by a highly demanding 30-min cognitive task, impaired repeated directional sprint and jumping ability. Neither repeated linear running nor singular CMJ were influenced by MF. Our findings may help to resolve the mixed findings in the extant literature. Specifically, strength endurance performance is vulnerable to MF when the task requires a higher level of cognitive engagement/demand. In sum, ecologically-valid studies on the effects of MF on strength endurance performance in sport would do well to consider the complexity and multiple determinants of athletic performance.

PRACTICAL APPLICATIONS

- Mentally demanding and fatiguing cognitive tasks may impair subsequent repetitive sprinting and jumping ability. Given that MF can also be caused by situational stressors and activities such as the use of smartphones, coaches and practitioners of sports with repetitive sprinting and jumping components (such as volleyball, football and basketball) should implement strategies to mitigate potential MF.
- To avoid potential negative effect of MF, coaches and practitioners could introduce cognitive recovery strategies for athletes (short-term solution), such as power napping, self-talk, and caffeine.
- Coaches and practitioners could use brain endurance training, a form of fatigue-inoculation training, to increase mental fatigue resilience (long-term solution).

REFERENCES

1. Van Cutsem J, Marcora S, De Pauw K, Bailey S, Meeusen R, Roelands B. The Effects of Mental Fatigue on Physical Performance: A Systematic Review. *Sports Med.* 2017;47(8):1569-1588. doi:10.1007/s40279-016-0672-0
2. Staiano W, Bosio A, Piazza G, Romagnoli M, Invernizzi PL. Kayaking performance is altered in mentally fatigued young elite athletes. *J Sports Med Phys Fitness.* 2019;59:1253–62. <https://doi.org/10.23736/S0022-4707.18.09051-5>
3. Habay J, Van Cutsem J, Verschueren J, et al. Mental Fatigue and Sport-Specific Psychomotor Performance: A Systematic Review. *Sports Med.* 2021;51(7):1527-1548. doi:10.1007/s40279-021-01429-6
4. Martin K, Thompson KG, Keegan R, Ball N, Rattray B. Mental fatigue does not affect maximal anaerobic exercise performance. *Eur J Appl Physiol.* 2015;115(4):715-725. doi:10.1007/s00421-014-3052-1
5. Fortes LS, de Lima-Júnior D, Fonseca FS, Albuquerque MR, Ferreira MEC. Effect of mental fatigue on mean propulsive velocity, countermovement jump, and 100-m and 200-m dash performance in male college sprinters. *Applied Neuropsychology: Adult.* 2021;0(0):1-10. doi:10.1080/23279095.2021.2020791
6. Kosack MH, Staiano W, Folino R, Hansen MF, Lombro S. The Acute Effect of Mental Fatigue on Badminton Performance in Elite Players. *Int J Sports Physiol Perform.* 2020;15(5):632-638. doi:10.1123/ijsp.2019-0361
7. Angius L, Merlini M, Hopker J, et al. Physical and Mental Fatigue Reduce Psychomotor Vigilance in Professional Football Players. *Int J Sports Physiol Perform.* Published online April 27, 2022:1-8. doi:10.1123/ijsp.2021-0387
8. Pageaux B, Marcora SM, Lepoint R. Prolonged mental exertion does not alter neuromuscular function of the knee extensors. *Med Sci Sports Exerc.* 2013;45(12):2254-2264. doi:10.1249/MSS.0b013e31829b504a
9. Alix-Fages C, Grgic J, Jiménez-Martínez P, Baz-Valle E, Balsalobre-Fernández C. Effects of Mental Fatigue on Strength Endurance: A Systematic Review and Meta-Analysis [published online ahead of print, 2022 Dec 12]. *Motor Control.* 2022;1-20. doi:10.1123/mc.2022-0051
10. Queiros VS de, Dantas M, Fortes L de S, et al. Mental Fatigue Reduces Training Volume in Resistance Exercise: A Cross-Over and Randomized Study. *Percept Mot Skills.* 2021;128(1):409-423. doi:10.1177/0031512520958935
11. Fortes LS, Lima-Júnior D, Barbosa BT, Faro HKC, Ferreira MEC, Almeida SS. Effect of mental fatigue on decision-making skill and visual search behaviour in basketball players: an experimental and randomised study. *Int J Sport Exerc Psychol.* Published online March 31, 2022. doi:10.1080/1612197X.2022.2058055
12. Pitts J, Bhatt T. Effects of mentally induced fatigue on balance control: a systematic review. *Exp Brain Res.* 2023;241(1):13-30. doi:10.1007/s00221-022-06464-x
13. Habay J, Uyenbroeck R, Van Droogenbroeck R, De Wachter J, Proost M, Tassignon B, De Pauw K, Meeusen R, Pattyn N, Van Cutsem J, Roelands B. Interindividual Variability in Mental Fatigue-Related Impairments in Endurance Performance: A Systematic Review and Multiple Meta-regression. *Sports Med Open.* 2023 Feb 20;9(1):14. doi: 10.1186/s40798-023-00559-7. PMID: 36808018; PMCID: PMC9941412
14. Mohr M, Krstrup P. Heat stress impairs repeated jump ability after competitive elite soccer games. *J Strength Cond Res.* 2013;27(3):683-689. doi:10.1097/JSC.0b013e31825c3266

15. Staiano W, Merlini M, Romagnoli M, Kirk U, Ring C, Marcora S. Brain Endurance Training Improves Physical, Cognitive, and Multitasking Performance in Professional Football Players. *Int J Sports Physiol Perform*. 2022 Nov 11;17(12):1732-1740. doi: 10.1123/ijsp.2022-0144. PMID: 36370703.
16. Tomporowski PD. Effects of acute bouts of exercise on cognition. *Acta Psychol (Amst)*. 2003;112(3):297-324. doi:10.1016/s0001-6918(02)00134-8
17. McKay AKA, Stellingwerff T, Smith ES, et al. Defining Training and Performance Caliber: A Participant Classification Framework. *Int J Sports Physiol Perform*. 2022;17(2):317-331. doi:10.1123/ijsp.2021-0451
18. Martin V, Sanchez-Sanchez J, Ramírez-Campillo R, Nakamura FY, Gonzalo-Skok O. Validity of the RSA-RANDOM test for young soccer players. *Int J Sports Med*. 2018;39(11):813–821. doi:10.1055/ a-0637-2094
19. Basner M, Dinges DF. Maximizing sensitivity of the Psychomotor Vigilance Test (PVT) to sleep loss. *Sleep*. 2011;34(5):581–591. doi:10.1093/sleep/34.5.581
20. Hart SG, Staveland LE. Development of NASA-TLX (Task Load Index): Results of empirical and theoretical research. In P. A. Hancock & N. Meshkati (Eds.), *Human mental workload* (pp. 139–183) 1988; North-Holland.
21. Girard O, Mendez-Villanueva A, Bishop D. Repeated-sprint ability - part I: factors contributing to fatigue. *Sports Med*. 2011;41(8):673-694. doi:10.2165/11590550-000000000-00000.
22. Borg G. Perceived exertion as an indicator of somatic stress. *Scand J Rehabil Med*. 1970;2:92–8. pmid:5523831.
23. Grgic J, Mikulic I, Mikulic P. Negative Effects of Mental Fatigue on Performance in the Yo-Yo Test, Loughborough Soccer Passing and Shooting Tests: A Meta-Analysis. *J Funct Morphol Kinesiol*. 2022;7(1):10. Published 2022 Jan 13. doi:10.3390/jfmk7010010
24. Verschueren J, Tassignon P, Proost M, Teugels A, Van Cutsem J, Roelands B, Verhagen E, Meeusen R (2020) Does mental fatigue negatively affect outcomes of functional performance tests? *Med Sci Sports Exerc* 52:2002–2010. <https://doi.org/10.1249/nss.0000000000002323>
25. Gantois P, Lima-Júnior L de, Fortes L de S, Batista GR, Nakamura FY, Fonseca F de S. Mental Fatigue From Smartphone Use Reduces Volume-Load in Resistance Training: A Randomized, Single-Blinded Cross-Over Study. *Percept Mot Skills*. 2021;128(4):1640–1659. doi:10.1177/00315125211016233
26. McMorris T, Hale BJ. Differential effects of differing intensities of acute exercise on speed and accuracy of cognition: a meta-analytical investigation. *Brain Cogn*. 2012;80(3):338-351. doi:10.1016/j.bandc.2012.09.001
27. Staiano W, Bonet LRS, Romagnoli M, Ring C. Mental Fatigue: The Cost of Cognitive Loading on Weight Lifting, Resistance Training, and Cycling Performance. *Int J Sports Physiol Perform*. 2023 Mar 9;18(5):465-473. doi: 10.1123/ijsp.2022-0356
28. Axelsen, J.L., Kirk, U. & Staiano, W. On-the-Spot Binaural Beats and Mindfulness Reduces the Effect of Mental Fatigue. *J Cogn Enhanc* 4, 31–39 (2020). doi.org/10.1007/s41465-019-00162-3
29. Proost M, Habay J, De Wachter J, et al. How to Tackle Mental Fatigue: A Systematic Review of Potential Countermeasures and Their Underlying Mechanisms. *Sports Med*. 2022;52(9):2129-2158. doi:10.1007/s40279-022-01678-z
30. Staiano W, Marcora M, Romagnoli M, et al. Brain Endurance Training improves endurance and cognitive performance in road cyclists, *Journal of Science and Medicine in Sport*, 2023; doi.org/10.1016/j.jsams.2023.05.008

FIGURES CAPTIONS

Figure 1. Schematic of the experimental protocol

Overall view of the experimental protocol. CMJ indicates counter movement jump test; PVT: psychomotor vigilant task; EC: Experimental condition; RSA: repeated-sprint ability random test; RJA: repeated-jump ability test; M-Vas: Visual Analogue mental fatigue scale; NASA TLX: National Aeronautic and Space Administration Task Load Index.

Figure 2. Cognitive and Physical Performance

2A Peak and Mean Time, during the Linear and Directional Phase of the Repeated Sprint Ability (RSA) Random Test for the experimental conditions: Mental fatigue and Control.

2B Peak Height of the Counter Movement Jump Test (CMJ) pre and post the experimental conditions on the left; Peak and Mean Height during the Repeated Jumping Ability (RJA) Test on the right; for the experimental conditions: Mental fatigue and Control.

2C Decrement Score expressed in percentage during the Repeated Sprint Ability (RSA) Random Test and the Repeated Jumping Ability (RJA) Test for the experimental conditions: Mental fatigue and Control.

2D Reaction Time (ms) of the Psychomotor Vigilance Test (PVT) before the experimental conditions (PRE EC), following the experimental conditions (POST EC) and following the physical tests (POST RSA-RJA) for the experimental conditions: Mental fatigue and Control.

Data are presented as mean (SD). \$ = significant condition x time interaction. # = significant main effect of condition. + = significant different from Control (simple main effect of condition).

Figure 3. Psycho-physiological measures

3A Heart Rate measured during the 12 Sprints in the Repeated Sprint Ability Random Test for the experimental conditions: Mental fatigue and Control.

3B Blood Lactate concentration measured before the experimental conditions (PRE EC), before the Repeated Sprint Ability Random Test (PRE RSA) and before the Repeated Jump Ability Test (PRE RJA) for the experimental conditions: Mental fatigue and Control.

3C RPE measured at completion of the 12 Sprints in the Repeated Sprint Ability Random Test for the experimental conditions: Mental fatigue and Control.

3D RPE measured at completion of the Repeated Jump Ability for the experimental conditions: Mental fatigue and Control.

Data are presented as mean (SD). \$ = significant condition x time interaction. * = significant main effect of time. # = significant main effect of condition.

Table 1: Subjective Ratings of Workload

	MENTAL DEMAND (0- 100)		PHYSICAL DEMAND (0- 100)		EFFORT (0- 100)		FRUSTRATION (0-100)	
	MF	CON	MF	CON	MF	CON	MF	CON
POST EC	62 (15)	35 (12)*	5 (3)	7 (3)	51(23)	19 (19)*	31 (19)	21 (25)
POST RSA- RJA	87 (13)	69 (16)*	89 (11)	86 (13)	91 (12)	78 (16)*	69 (21)	50 (17)*

Abbreviations: EC = Experimental condition, MF = Mental fatigue condition, CON = Control condition

Note: Data are presented as mean (SD), * significant difference between conditions ($p < .05$)

CRediT Author Statement

	Staiano W	Lluís Salazar	Romagnoli M	Ring C
Conceptualization	X	X		X
Methodology	X		X	
Software	X	X		
Validation	X	X		
Formal analysis	X			X
Investigation	X	X		
Resources	X		X	X
Data Curation	X		X	
Writing - Original Draft	X			X
Writing - Review & Editing	X	X	X	X
Visualization	X	X		
Supervision	X			X
Project administration	X		X	
Funding acquisition				

We would like to confirm here below the contribution of each authors using the typical 14 role-model.

I, hereby, acknowledge that all authors have revised and agreed upon the contribution table here presented.

Acknowledgments

We would like to thank all the participants who took part in the study.

Funding Information

The studies did not receive any funding

Declaration of Interest

We declare that any author the present study does not have any conflict or personal interest related to the data collected.

Confirmation of Ethical Compliance

The study was approved by the Ethics Committee for the University of Valencia in accordance with the Declaration of Helsinki.




Figure 1




Figure 2







Figure 3