



Changes in alcohol consumption during the COVID-19 among first-year university students in Spain, considering the risk of problematic use – UniHcos project



Lorena Botella-Juan ^{a, b}, María Morales-Suárez-Varela ^{c, d}, Carmen Amezcua-Prieto ^{c, e, f}, Ramona Mateos-Campos ^g, Carlos Ayán-Pérez ^h, Antonio José Molina ^{a, b, c, *}, Rocío Ortiz-Moncada ⁱ, Ana Delgado-Parrilla ^j, Gemma Blázquez-Abellán ^k, Miguel Delgado-Rodríguez ^{c, l}, Jessica Alonso-Molero ^m, Tania Fernández-Villa ^{a, b, c}

^a Department of Biomedical Sciences, Area of Preventive Medicine and Public Health, Faculty of Health Sciences, Universidad de León, 24071 León, Spain

^b The Research Group in Gene-Environment and Health Interactions (GIIGAS), Institute of Biomedicine (IBIOMED), Universidad de León, 24071 León, Spain

^c Consortium for Biomedical Research in Epidemiology & Public Health (CIBER Epidemiología y Salud Pública-CIBERESP), 28029 Madrid, Spain

^d Department of Preventive Medicine and Public Health, Food Sciences, Toxicology and Legal Medicine, Universidad de Valencia, 46100 Burjassot, Spain

^e Department of Preventive Medicine and Public Health, Universidad de Granada, 18016 Granada, Spain

^f Instituto de Investigación Biosanitaria (ibs.Granada), 18014 Granada, Spain

^g Area of Preventive Medicine and Public Health, Department of Biomedical and Diagnostic Sciences, Universidad de Salamanca, 37007 Salamanca, Spain

^h Well-Move Research Group, Department of Special Didactics, Universidad de Vigo, 36310 Vigo, Spain

ⁱ Area of Preventive Medicine and Public Health, Food and Nutrition Research Group, Universidad de Alicante, 03550 Alicante, Spain

^j Centre for Research on Natural Resources, Health, and Environment (RENSMA), Universidad de Huelva, 21071 Huelva, Spain

^k Department of Medical Sciences, Faculty of Pharmacy, Universidad de Castilla-La Mancha, 02008 Albacete, Spain

^l Department of Health Sciences, Universidad de Jaén, 23071 Jaén, Spain

^m Department of Preventive Medicine and Public Health, Universidad de Cantabria-IDIVAL, 39008 Santander, Spain

ARTICLE INFO

Article history:

Received 5 March 2024

Received in revised form

23 May 2024

Accepted 21 June 2024

Keywords:

alcohol drinking in college

university student drinking

alcoholism

binge drinking

COVID-19

ABSTRACT

The aim of this study was to evaluate the possible impact of the COVID-19 pandemic on prevalence rates and self-reported changes in patterns of alcohol use among first-year university students in Spain, considering the risk of problematic alcohol use. A serial cross-sectional study based on the uniHcos project was carried out. Data from 10 518 first-year university students (73.3% female, mean age 19 (SD = 1.6)) collected between 2012 and 2022 were analysed. The evolution of the pooled prevalence rates during the time series was analysed and the risk of problematic alcohol consumption was assessed using the AUDIT. Also, self-reported changes in alcohol use patterns during the pandemic were assessed. According to the results, during the COVID-19 pandemic, the prevalence of alcohol use in the past 30-days was reduced (76.3% in 2019 vs. 63.7% in COVID-19) increasing again in the New Normal period. Thus, a similar pattern in the practice of binge drinking was observed. Regarding the AUDIT score, 21.7% (95%CI 20.9, 22.6) of the students had harmful alcohol consumption, with a higher proportion among males. In the multivariable logistic models, a higher AUDIT score was significantly associated (p-value < 0.001) with being male and living with roommates. According to self-reported changes in consumption patterns during the COVID-19 pandemic, a higher proportion of participants with harmful use reported an increase in alcohol consumption compared to those at low-risk (43% vs 19%). Finally, despite the overall reduction in drinking prevalence during COVID-19, changes were not equal for all students and depended on their previous level of problematic drinking, highlighting that this should be considered in the development of strategies against alcohol use in this population.

© 2024 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

* Corresponding author. Department of Biomedical Sciences, Area of Preventive Medicine and Public Health, Faculty of Health Sciences, Universidad de León, 24071 León, Spain.

E-mail address: ajmolt@unileon.es (A.J. Molina).

Introduction

The onset of the COVID-19 pandemic in 2020 led to the imposition of strong restrictive measures to prevent its spread worldwide. In accordance with the legislation ([Real Decreto 463/2020](#); [Real Decreto 926/2020](#)) these measures in Spain were divided into two phases. First, the initial phase, known as the “first-lockdown”, which covered the first months of the pandemic (March–June 2020), characterised by the stress and anxiety of the novelty, mandatory home lockdown, lack of knowledge, fear, and the beginning of the use of masks. Then, in the second phase (July 2020–May 2021) the measures were lighter, however, there were still curfews, social distancing and the use of masks. After this period, from June 2021, the return to normality began (New Normal period).

These changes could significantly impact the population's lifestyle habits, such as substance use patterns ([Schmidt et al., 2021](#)). About this, the changes in alcohol use during the COVID-19 pandemic are of great interest with a particular focus on the university population and especially on the first-year university students. The university stage is a turning point in students' lives as it coincides with the transition into adulthood, greater independence, and different social circles ([Locke et al., 2015](#)). Additionally, the start of this stage frequently coincides with the initiation of substance use experimentation ([El Ansari et al., 2020](#)) and with the increase in alcohol consumption and risky drinking patterns ([Flaudias et al., 2021](#)). Furthermore, the transition from the family home to independent living could be a significant risk factor for alcohol consumption, due to the influence of roommates and the sense of freedom ([Moure-Rodríguez et al., 2020](#)).

The *European Health Interview Survey* (EHIS) indicates that Spain is the second most frequent alcohol consumer in Europe ([Eurostat, 2021](#)). Among young people in Spain, the *National Survey on Alcohol and other Drugs in Spain* (EDADES) ([OEDA, 2022](#)) reported that 62.7% of people aged 15–24 drank alcohol in the past month. Although the legal age for the purchase and consumption of alcoholic drinks in Spain is 18, the average age of onset is 16.5 years. Consequently, by the time students enrol in university, many of them have already engaged in the consumption of alcohol. In addition, this population has been identified as one of the populations with the highest rates of alcohol consumption worldwide ([White & Hingson, 2014](#); [Yockey et al., 2020](#)). Early alcohol consumption has been linked to an increased risk for later alcohol dependence, mental and social problems ([Isaksson et al., 2019](#)) and the development of severe physical diseases ([Messina et al., 2021](#)). Additionally, binge drinking, or heavy episodic drinking, is prevalent among college students ([Castañeda et al., 2021](#)), and has been linked to a multitude of adverse outcomes ([Valencia Martín et al., 2020](#)). The typical motivations for alcohol consumption among college students include greater social integration, having a good time and a low-risk perception ([Mereu et al., 2021](#)). Hazardous drinkers, however, have reasons such as enhancement or coping with negative emotions, social anxiety, depression, alcohol dependence and poor mental health ([Bresin & Mekawi, 2021](#); [Messina et al., 2021](#)).

The onset of the COVID-19 pandemic has prompted concerns regarding the potential impact on alcohol use patterns. Social events were significantly reduced, yet alcohol sales remained accessible in Spain. Concurrently, emotional distress and uncertainty increased, representing a risk factor for alcohol use ([Clay & Parker, 2020](#)). In this context, some authors have observed an increase in alcohol consumption among university students ([Busse et al., 2021](#); [Charles et al., 2021](#)), although the majority of studies have reported a decrease in the overall prevalence of consumption ([Babicki, 2022](#); [Bollen et al., 2021](#); [Evans et al., 2021](#); [Gesualdo &](#)

[Pinquart, 2021](#); [Heradstveit et al., 2022](#); [Tavolacci et al., 2021](#); [van Hooijdonk et al., 2022](#)), mainly attributed to less availability and the reduction of social opportunities ([Evans et al., 2021](#); [Gesualdo & Pinquart, 2021](#)). However, the majority of these studies have only considered consumption during the first-lockdown period and have treated this population as a single entity, without differentiating between drinkers' profiles.

Moreover, the research on this topic in Spain is limited. Most studies do not focus on alcohol consumption as the primary objective, but rather on broader changes in health habits, such as physical activity, diet, and the impact of the pandemic on mental health ([Lázaro-Pérez et al., 2020](#); [Millán-Jiménez et al., 2021](#); [Romero-Blanco et al., 2020](#); [Ruiz-Zaldibar et al., 2022](#); [Zapata et al., 2022](#)).

A comprehensive examination of the changes in alcohol consumption among first-year university students in Spain is worthy of consideration. Such an investigation could provide insights into how these young individuals respond to unique circumstances and the factors that may influence their patterns of alcohol consumption. Furthermore, the identification of the different profiles of drinkers in the university community would contribute to the design of appropriate strategies and interventions to deal with the problem of alcohol consumption considering the perspective of the COVID-19 pandemic.

In light of the above, the objective of this study was to fill the gap on the effect that the COVID-19 pandemic and the measures imposed may have had on the prevalence and patterns of alcohol consumption among first-year university students in Spain, compared to the previous trends and considering the risk of problematic drinking and the differences by sex.

Methods

Participants and data collection

A serial cross-sectional study based on the basal questionnaire of the uniHcos project was conducted ([Fernández-Villa et al., 2013](#)). The study population were 11485 First-Year university students from eleven Spanish universities who participated between 2012 and May 2022. The uniHcos questionnaire is online and self-administered, with 471 assessable items about lifestyle habits. Since 2020, questions designed to evaluate the impact of the COVID-19 pandemic on these habits were included.

The students were recruited through their university account email. The invitation included information about the project objectives, data use, confidentiality mandatory informed consent and the survey link in the SphinxOnline® platform (v. 4.26, Le Sphinx Développement, Chavanod, France). The procedure was conducted in accordance with the current data protection regulations and the Code of Ethics of the World Medical Association (Declaration of Helsinki). All ethics committees of the collaborating universities evaluated and approved the project. The students participated voluntarily, without any kind of reward.

For the purposes of analysis, the age range was limited to 17–24 years. Of the initial sample, 973 participants (8.5%) were excluded due to being older than the age indicated. This decision was made in order to ensure that the population was as homogeneous as possible and represented the majority of first-year university students in Spain. The final sample consisted of 10 518 students, 73.3% female, with a mean age (standard deviation) of 19.0 (1.6) years.

The data was classified into three periods as follows: from 2012 to 2019 the surveys were grouped by the year of filling, considering this period as Before COVID-19 period. The second period established was COVID-19 pandemic and it included from the imposition

of the “Alarm State”, a period of severe restrictions in Spain, until its repeal (March 2020 to May 2021) according to the legislation (Real Decreto 463/2020; Real Decreto 926/2020). The third period was the New Normal period and it included from June 2021 to May 2022 and is the period of return to normality in most aspects.

In order to analyse changes in self-reported alcohol consumption patterns during the COVID-19 pandemic, this period was divided into two sub-periods: “COVID-19 pandemic (first-lockdown)” (March 2020–June 2020) and “COVID-19 pandemic (not first-lockdown)” (July 2020–May 2021).

Study variables

The prevalence of alcohol consumption at different frequencies was explored, as well as the practice of binge drinking (BD), defined as the consumption of 5 or more drinks for men and 4 or more drinks for women on a single occasion within approximately 2 h (Courtney & Polich, 2009; OEDA, 2022; Vasconcelos et al., 2021). Furthermore, the risk of problematic alcohol use was evaluated using the *Alcohol Use Disorders Identification Test* (AUDIT) (Saunders et al., 1993). This test has been validated for its use in university populations and in an online format (Ballester et al., 2021; Kokotailo et al., 2004). For the analysis two categories were defined: “low risk of problematic use” (score 0–7) and “medium/high risk (hazardous/harmful use)” (score ≥ 8). For some analysis, the latter category was further subdivided into “medium risk” (score 8–15) and “high risk” (score ≥ 16), in line with other authors (Killgore et al., 2021; WHO et al., 2001). Covariables such as sex, age and cohabitants were considered.

To investigate the self-perceived changes in drinking patterns during the COVID-19, since the survey sent out in 2020–2021 two questions were included on this topic. One concerning changes during the first-lockdown (COVID-19 pandemic first-lockdown): “During the first-lockdown period, how your alcohol use was in comparison with before the first-lockdown?”, and one about the perception throughout the pandemic period (COVID-19 pandemic not first-lockdown): “Since the arrival of COVID-19 in Spain, how was your alcohol use?”, with possible answers of both questions being (1) “It has increased”, (2) “The use is the same”, (3) “The use has decreased”, (4) “I’m not an alcohol user”.

Data analysis

The pooled prevalence rates of alcohol use and binge drinking for each period and their 95% Confidence Interval (95%CI) were analysed. To identify periods with significant changes in the prevalence trends over the time series, joinpoint regressions models were estimated calculating Annual Percent Changes (APCs). Up to three joinpoints were allowed, and p-values less than 0.05 were considered statistically significant. The “pairwise comparison” option within the software was used to assess whether trends differed significantly between sexes. Furthermore, bivariate and multivariable adjusted logistic regressions were performed to assess the association between the AUDIT score and different variables. A descriptive analysis of the self-reported patterns of alcohol use during the COVID-19 pandemic was also conducted.

To assess statistically significant differences between categorical variables the Pearson's Chi-Square analysis was used. The statistical significance was established at 95% ($\alpha = 0.05$). All analyses were performed using the software STATA 16 (StataCorp., 2019), except for joinpoint regressions that were developed using the *Joinpoint Regression Program*, Version 5.1.0.0 (Statistical Research and Applications Branch, National Cancer Institute).

Results

Table 1 shows the socio-demographic characteristics of the sample. Most participants were female (73%) and most of the students used to live with their family (44%) or with roommates/friends (45%).

Evolution of prevalence of alcohol use

According to Fig. 1, throughout the time series, the prevalence of alcohol use ever in lifetime was very high reaching its highest value in 2017 (94.7%). About the past 12-months, the prevalence rates remained stable although the lowest value was found in the COVID-19 period (87.9%). Regarding the past 30-days, a large decrease was observed in the COVID-19 period, where the prevalence dropped to 63.7%, increasing again in the New Normal period (77.8%). A similar pattern was observed with regard to the prevalence of binge drinking in the past 30-days (46.0% in 2019 to 31.3% in COVID-19 and 44.4% in the New Normal).

The results of joinpoint regression models supported these changes, with trend change points identified in COVID-19 and New Normal periods for the prevalence of past 30-days ($APC_{2018-2019} = -8.3/APC_{COVID-19-NewNormal} = 19.0$) and binge drinking ($APC_{2018-2019} = -17.2/APC_{COVID-19-NewNormal} = 32.3$), with no significant sex differences observed. In contrast, lifetime and past 12-months prevalence rates showed no change due to the effect of the pandemic, with small trend changes not exceeding an APC of 2.5%. The complete analysis and Annual Percent Changes (APCs) are available in the *Supplementary Material 1*.

Differences in alcohol use prevalences and associated factors according to AUDIT score

Table 2 shows that 21.7% of the total sample scored ≥ 8 points in AUDIT with statistically significant differences by sex ($p < 0.001$). In terms of the study period, during the COVID-19 period the percentage of students in the medium/high-risk level decreased from 22.0% in before COVID-19 to 17.7%, increasing again in the New Normal period. Also, according to the multivariable logistic regression model (Table 2), an association was found between being male ($aOR = 1.49$; p -value < 0.001) and live with roommates ($aOR = 1.69$, p -value < 0.001) and an increased risk of a positive AUDIT screening, while the COVID-19 period was a protective factor ($aOR = 0.74$, p -value = 0.007). In bivariate analysis, binge drinking practice in the past 30 days ($OR = 10.06$) were statistically significant associated ($p < 0.001$) with an AUDIT score of ≥ 8 points (data not shown).

Table 1
Sociodemographic characteristics of the sample (first-year university students), collected between 2012 and 2022 in Spain.

	n (N = 10 518)	% (95%CI)
Sex		
Female	7710	73.3 (72.5, 74.1)
Male	2808	26.7 (25.9, 27.5)
Period of study		
Before COVID-19	9331	88.7 (88.1, 89.3)
COVID-19 pandemic	619	5.9 (5.4, 6.3)
New Normal	568	5.4 (5.0, 5.8)
People the student lives with		
Family	4614	43.9 (42.9, 44.8)
Roommates/Friends	4688	44.6 (43.6, 45.5)
Alone	856	8.1 (7.6, 8.7)
Other	360	3.4 (3.1, 3.8)

Note. Data collection of the periods: Before COVID-19 period: 2012–2019, COVID-19 pandemic: March 2020–May 2021, New Normal period: June 2021–May 2022.




Fig. 1. Evolution, from 2012 to 2022, of the prevalence (%) of alcohol use in different frequencies: ever, past 12-months, past 30-days and binge drinking (past 30-days) in first-year university students in Spain. Error bars show the 95%CI.

Upon disaggregating the data on the prevalence of alcohol consumption in the past 30 days by AUDIT score, it was found that the decrease observed during the period of the COVID-19 pandemic was not as pronounced among those with an AUDIT ≥ 8 (97.2% before COVID-19 to 89.1% during COVID-19) as among those with a low-risk level (73.3%–58.3%) (Supplementary Material 2).

Changes in the self-reported patterns of use during first-lockdown and all COVID-19 pandemic, by AUDIT score

According to our findings, of the students who participated after the pandemic ($n = 1187$) 63.8% reported drinking during the first-

lockdown. Among them, 55.9% reported a decrease in their consumption, while 28.1% maintained it and 15.9% increased it. Regarding the changes during all COVID-19 pandemic (not first-lockdown), 72.6% declared alcohol use. Of them, 35.9% decreased their alcohol use, while 38.1% maintained it and 25.9% increased it.

Stratifying the results according the AUDIT score (Fig. 2), during the first-lockdown, the 26.9% of medium/high-risk students reported an increase in alcohol consumption, compared to 11.8% of low-risk students ($p < 0.001$). Similar results were observed in all COVID-19 period. 43.0% of the participants with harmful use increased their consumption compared to 19.8% of those at low-risk ($p < 0.001$).

Table 2

Prevalence, bivariate and multivariable logistic regressions for the association of risk of problematic alcohol use (assessed with AUDIT) and different variables. Data from a cross-sectional sample of first-year Spanish university students.

AUDIT SCORE ≥ 8 points							
Prevalence	% (95%CI)	Bivariate			Multivariable		
		OR	95%CI	p-value	aOR	95%CI	p-value
Total	21.7 (20.9, 22.6)	—			—		
Sex							
Female	19.5 (18.7, 20.4)						
Male	27.9 (26.3, 29.6)	1.59	1.44, 1.75	<0.001	1.65	1.49, 1.82	<0.001
Period of Study							
Before COVID-19	22.0 (21.2, 22.9)						
COVID-19 pandemic	17.7 (14.9, 20.9)	0.76	0.62, 0.95	0.013	0.74	0.59, 0.92	0.007
New Normal	22.0 (18.8, 25.6)	0.99	0.81, 1.22	0.993	0.92	0.74, 1.14	0.448
People they live with							
Parents	17.0 (16.0, 18.1)						
Roommates/Friends	26.8 (25.6, 28.2)	1.66	1.51, 1.83	<0.001	1.69	1.53, 1.87	<0.001
Age (ref=17)^a		0.99	0.96, 1.03	0.879	0.99	0.96, 1.02	0.592

Note. Data collection of the periods: Before COVID-19 period: 2012–2019, COVID-19 pandemic: March 2020–May 2021, New Normal period: June 2021–May 2022. Adjusted logistic regression models by sex, age, period of study and university of precedence.

^a Mean age (SD) = 19.0 (1.6) years.

For all COVID-19 pandemic, we stratified the analysis in three AUDIT score levels (low, medium, and high, data not shown in the graph) founding that 75.9% (CI95% 57.0, 88.2) of participants at high-risk increased their consumption, 13.8% (CI95% 5.2, 31.9) maintained it and only 10.3% (CI95% 3.3, 28.0) decreased it.

Changes in the self-reported patterns of use during first-lockdown and all COVID-19 pandemic, by sex

As observed in Fig. 3, during the COVID-19 (first-lockdown) there was a higher proportion of females reporting reducing their alcohol consumption than males, with statistically significant differences ($p < 0.001$). Regarding the changes during all COVID-19 pandemic (not first-lockdown), there was not statistically significant differences by sex in the change of patterns of use ($p = 0.118$).

Discussion

The main objective of this study was to explore the possible impact of the COVID-19 pandemic on alcohol consumption among first-year university students in Spain, considering risky drinking and with a 10-year evolutionary perspective. The results indicated that throughout the time series, the lifetime prevalence of alcohol consumption was above 95%, while the prevalence rate in the past 12-months was close to 90%. Both prevalence rates remained stable over time.

However, a significant decrease was observed in the past 30-days frequency of use during the COVID-19 period (76.3% in 2019 vs 63.7% in the COVID-19), with a subsequent increase in the New

Normal period. This pronounced decline in alcohol consumption is consistent with prior studies on its use during periods of high restrictions (Del Valle-Vera et al., 2021; Kilian et al., 2022; Manthey et al., 2021; OEDA, 2021). Indeed, the outbreak of the COVID-19 was identified as a protective factor against alcohol consumption, according to other authors (Ruiz-Zaldibar et al., 2022).

Nevertheless, the return to previous levels in the New Normal may indicate that the change was a one-off process and that alcohol consumption is closely related to social context. This behaviour has been observed in other studies (Blumberg et al., 2023). This is also supported by other authors who have identified that the main reason for the decline in alcohol consumption among university students was the reduction in social interaction opportunities (Bramness et al., 2021; Fruehwirth et al., 2021; Jackson et al., 2021). Since this paper presents a 10-year perspective on alcohol use prevalence, the sharp decrease in the past-30 days frequency during the COVID-19 period is of interest. This contrasts with the other frequencies of use, which have remained stable, thus supporting the conclusion that these observed changes can indeed be attributed to the outbreak of the pandemic.

Furthermore, the same phenomenon has been observed in the reduction in binge drinking during the COVID-19 period, which also increased again in the new normal period. This has been observed in similar populations (Bonar et al., 2021; Fruehwirth et al., 2021; Tavolacci et al., 2021), probably because of its character of “context-related habit” too (Vasconcelos et al., 2021). It should be noted that the reduction in this practice has been associated with a reduction in the overall prevalence (Kilian et al., 2021).




Fig. 2. Changes in drinking patterns of first-year Spanish university students during COVID-19 pandemic (first-lockdown) (March 2020–June 2020) and COVID-19 pandemic (not first-lockdown) (July 2020–May 2021), by AUDIT score (cut-off point ≥ 8). Error bars show the 95%CI. *Show statistically significant differences in patterns.




Fig. 3. Changes in drinking patterns of first-year Spanish university students during COVID-19 pandemic (first-lockdown) (March 2020–June 2020) and COVID-19 pandemic (not first-lockdown) (July 2020–May 2021), by sex. Error bars show the 95%CI. *Show statistically significant differences in patterns.

On another note, our results highlight that within the university population, the changes during COVID-19 could be related to the profile of the drinker. The percentage of students with medium/high-risk drinking decreased during the COVID-19, increasing again in the New Normal period, as found by other authors (Heradstveit et al., 2022; Kosendiak et al., 2022). The use of the AUDIT questionnaire may explain this, as it refers in four out of ten questions to frequency or quantity consumed, and both parameters were reduced during COVID-19 (Blumberg et al., 2023; Del Valle-Vera et al., 2021; Jaffe et al., 2021).

Despite these results, we observed differences in the pattern of use during COVID-19 pandemic according to the score obtained in AUDIT. A higher proportion of participants in the medium/high-risk category increased or maintained their consumption than in the low-risk category. Moreover, this difference was even more pronounced in the segregated analysis of those at high risk, with various authors also identifying this phenomenon (Chodkiewicz et al., 2020; Matone et al., 2022; Rossow et al., 2021; Schmidt et al., 2021; Wysokińska & Kotota, 2022). This result indicates that when confronted with the similar social changes and stressors, alcohol users do not respond in the same way, and they probably change their consumption in accordance with their previous risk of problematic use. Furthermore, residing away from family was found as a risk factor for a positive AUDIT screening. This finding has been reported by other authors (van Hooijdonk et al., 2022; White et al., 2020; Wicki et al., 2010; Zysset et al., 2022), indicating that the co-living situation should be considered.

It is worth noting that the existing literature has shown strong correlation between an increased risk of problematic use and impaired mental health, or mental health disorders (Chang et al.,

2021; Charles et al., 2021; Gavurova et al., 2021; Kenney et al., 2018; Pudddephatt et al., 2021), so the strong evidence of worsening mental health among students during COVID-19 (Gogoi et al., 2022; Padrón et al., 2021; Yuan & You, 2022), may support our results.

In addition, no significant differences in the prevalence of alcohol use by sex were found and while variations in patterns were observed during the COVID-19 first-lockdown, they were not in the all COVID-19 pandemic period. The results are in alignment with the current convergence of consumption patterns between men and women, which is attributed to new changes in women's lives. These changes are leading women to consume alcohol in the same way as men in formally egalitarian societies (Holmila & Raitasalo, 2005; Iwamoto & Mui, 2020). However, there were significant differences in risky consumption, with a higher proportion of males. Also, being male was found as a risk factor for problematic drinking, a fact that has been described in the current literature (Aresi et al., 2022; Babicki, 2022; Steffen et al., 2021; Zysset et al., 2022). Nevertheless, despite the narrowing of the overall prevalence by sex, it seems that there are still differences in the risk of problematic use (López-Moreno et al., 2021), which is relevant for future studies.

The results of this study point to the need to implement targeted strategies for university students, given their specific drinking profiles, as not all drinkers have the same patterns. In light of this, it is important to note that strategies aimed at reducing the availability of alcohol on campus may be useful for those at low risk, as it has been suggested that reduced availability may have reduced consumption during COVID-19 (Kilian et al., 2021). However, these strategies may not be sufficient for those who have already

developed a high risk of alcohol consumption, leaving out a large group of students.

For this reason, it would be beneficial to implement strategies such as the one proposed by Benjet et al. (2022) within the Spanish university community. This strategy involves the administration of a brief online survey, which then allows an algorithm to determine each student's risk based on the AUDIT score and other relevant factors, such as those identified in this study (sex, cohabitation and patterns of use). This approach would enable the identification of at-risk first-year students, thereby preventing the emergence of significant future issues.

Previous research has demonstrated that alcohol consumption during the first year of university is predictive of subsequent drinking behaviour (Prince et al., 2019). The capacity to promptly identify at-risk students would permit them to be aware of their level of risk and to be recommended for useful strategies for example face-to-face participation in Brief Alcohol Interventions (BAIs) with evidence (Hennessy et al., 2019), as well as assistance and mitigation guidelines, such as Protective Behavioural Strategies (PBS) (Linden-Carmichael et al., 2019).

Despite the meaningful findings provided, this study is not without limitations. First, the limitations of a cross-sectional study design prevent us from providing causality. Secondly, data collection was obtained by a self-administered questionnaire, so self-reported consumption data is a measure with restrictions in generalizability to other populations. Finally, this work also has limitations related to the sample, considering the difference in sample size in some periods and a higher proportion of women in the sample. However, more women are entering university in Spain (56.0%), especially in some fields like health sciences (71.4%) (Ministerio de Universidades, 2022), so it is hoped that this and the disaggregation of data by sex will control for a possible underestimation of men.

Conclusions

During COVID-19, a reduction in the overall prevalence of past 30-day alcohol consumption and binge drinking was observed. However, there were notable differences in the participants' self-reported changes of drinking patterns according to their previous risk of problem drinking. Thus, those students with a medium/high risk level increased their alcohol consumption to a greater extent during the pandemic. Being male and living without the family were associated with risky drinking. This study highlights that any strategy targeting the university population must consider not only the overall prevalence of alcohol consumption, but also the risk of problem drinking of the students, in order to provide tailored interventions that can be truly effective for students' needs.

Data availability statement

The datasets analyzed in the current study are available from the corresponding author upon reasonable request.

Ethics approval statement

The procedure was carried out following current data protection regulations and following the Code of Ethics of the World Medical Association (Declaration of Helsinki of 1975, as revised in 2000). All ethics committees of the collaborating universities evaluated and approved the project. Informed consent was obtained from all patients for being included in the study. The students participated voluntarily, without any kind of reward.

Funding

This research was funded by National Drug Plan from the Ministry of Health, Social Services and Equality of Spain: 2010|145, 2013|034, 2020|030; Instituto de Salud Carlos III: PI16/01947. Lorena Botella-Juan: predoctoral contract financed by Junta de Castilla y León—Consejería de Educación (Spain) using the European Social Fund (EDU/842/2022).

CRediT authorship contribution statement

Lorena Botella-Juan: Writing – review & editing, Writing – original draft, Visualization, Methodology, Investigation, Formal analysis, Data curation, Conceptualization. **María Morales-Suárez-Varela:** Writing – review & editing, Methodology, Formal analysis, Data curation, Conceptualization. **Carmen Amezcua-Prieto:** Writing – review & editing, Data curation, Conceptualization. **Ramona Mateos-Campos:** Writing – review & editing, Data curation, Conceptualization. **Carlos Ayán-Pérez:** Writing – review & editing, Data curation, Conceptualization. **Antonio José Molina:** Writing – review & editing, Visualization, Supervision, Methodology, Formal analysis, Data curation, Conceptualization. **Rocío Ortiz-Moncada:** Writing – review & editing, Data curation, Conceptualization. **Ana Delgado-Parrilla:** Writing – review & editing, Data curation, Conceptualization. **Gemma Blázquez-Abellán:** Writing – review & editing, Data curation, Conceptualization. **Miguel Delgado-Rodríguez:** Writing – review & editing, Data curation, Conceptualization. **Jessica Alonso-Molero:** Writing – review & editing, Data curation, Conceptualization. **Tania Fernández-Villa:** Writing – review & editing, Writing – original draft, Visualization, Supervision, Project administration, Methodology, Funding acquisition, Formal analysis, Data curation.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors acknowledge the help and collaboration of the students who participated in the study.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.alcohol.2024.06.008>.

References

- Aresi, G., Sorgente, A., Cleveland, M. J., & Marta, E. (2022). Patterns of alcohol use among Italian young adults before and during a COVID-19 lockdown: A latent class analysis study. *Journal of Prevention*, 43(2), 191–208. <https://doi.org/10.1007/S10935-022-00675-2>, 2022.
- Babicki, M. (2022). Use of alcohol, cannabinoids, psychostimulants, and sedatives before and during the COVID-19 pandemic among students in 40 European countries. *International Journal of Environmental Research and Public Health*, 19(22), 14879. <https://doi.org/10.3390/ijerph192214879>
- Ballester, L., Alayo, I., Vilagut, G., Almenara, J., Cebrià, A. I., Echeburúa, E., Gabilondo, A., Gili, M., Lagares, C., Piqueras, J. A., Roca, M., Soto-Sanz, V., Blasco, M. J., Castellví, P., Forero, C. G., Mortier, P., & Alonso, J. (2021). Validation of an online version of the alcohol use disorders identification test (AUDIT) for alcohol screening in Spanish university students. *International Journal of Environmental Research and Public Health*, 18(10), 5213. <https://doi.org/10.3390/IJERPH18105213>
- Benjet, C., Mortier, P., Kiekens, G., Ebert, D. D., Auerbach, R. P., Kessler, R. C., Cuijpers, P., Green, J. G., Nock, M. K., Demeyttenaere, K., Albor, Y., & Bruffaerts, R. (2022). A risk algorithm that predicts alcohol use disorders among college

- students. *European Child & Adolescent Psychiatry*, 31(7), 1–11. <https://doi.org/10.1007/s00787-020-01712-3>
- Blumberg, M. J., Lo, L. A., Harrison, G. W., Dodwell, A., Irwin, S. H., & Olmstead, M. C. (2023). Differential impacts of perceived social support on alcohol and cannabis use in young adults: Lessons from the COVID-19 pandemic. *University of Toronto Medical Journal*, 100(1), 28–39. <https://doi.org/10.33137/utmj.v100i1.38056>
- Bollen, Z., Pabst, A., Creupelandt, C., Fontesse, S., Lannoy, S., Pinon, N., & Muraige, P. (2021). Prior drinking motives predict alcohol consumption during the COVID-19 lockdown: A cross-sectional online survey among Belgian college students. *Addictive Behaviors*, 115, Article 106772. <https://doi.org/10.1016/j.addbeh.2020.106772>
- Bonar, E. E., Parks, M. J., Gunlicks-Stoessel, M., Lyden, G. R., Mehus, C. J., Morrell, N., & Patrick, M. E. (2021). Binge drinking before and after a COVID-19 campus closure among first-year college students. <https://doi.org/10.1016/j.addbeh.2021.106879>
- Bramness, J. G., Bye, E. K., Moan, I. S., & Rossow, I. (2021). Alcohol use during the COVID-19 pandemic: Self-reported changes and motives for change. *European Addiction Research*, 27(4), 257–262. <https://doi.org/10.1159/000515102>
- Bresin, K., & Mekawi, Y. (2021). The “why” of drinking matters: A meta-analysis of the association between drinking motives and drinking outcomes. *Alcoholism: Clinical and Experimental Research*, 45(1), 38–50. <https://doi.org/10.1111/ACER.14518>
- Busse, H., Buck, C., Stock, C., Zeeb, H., Pischke, C. R., Fialho, P. M. M., Wendt, C., & Helmer, S. M. (2021). Engagement in health risk behaviours before and during the covid-19 pandemic in German university students: Results of a cross-sectional study. *International Journal of Environmental Research and Public Health*, 18(4), 1–16. <https://doi.org/10.3390/ijerph18041410>
- Castañeda, G., Colby, S. E., Olfert, M. D., Barnett, T. E., Zhou, W., Engel, M. D., & Mathews, A. E. (2021). The role of alcohol-related proactive dietary restriction to feel the psychoactive effects of alcohol faster on binge drinking frequency among freshmen college students. *Substance Use & Misuse*, 56(9), 1266–1274. <https://doi.org/10.1080/10826084.2021.1914104>
- Chang, W.-P., Stuart, H., & Chen, S.-P. (2021). Alcohol use, Drug use, and psychological distress in university students: Findings from a Canadian campus survey. *International Journal of Mental Health and Addiction*, 20, 2366–2381. <https://doi.org/10.1007/s11469-021-00519-w>
- Charles, N. E., Strong, S. J., Burns, L. C., Bullerjahn, M. R., & Serafine, K. M. (2021). Increased mood disorder symptoms, perceived stress, and alcohol use among college students during the COVID-19 pandemic. *Psychiatry Research*, 296, Article 113706. <https://doi.org/10.1016/j.psychres.2021.113706>
- Chodkiewicz, J., Talarowska, M., Miniszewska, J., Nawrocka, N., & Bilinski, P. (2020). Alcohol consumption reported during the COVID-19 pandemic: The initial stage. *International Journal of Environmental Research and Public Health* 2020, 17(13), 4677. <https://doi.org/10.3390/ijerph17134677>. Vol. 17, Page 4677.
- Clay, J. M., & Parker, M. O. (2020). Alcohol use and misuse during the COVID-19 pandemic: A potential public health crisis? *The Lancet Public Health*, 5(5), e259. [https://doi.org/10.1016/S2468-2667\(20\)30088-8](https://doi.org/10.1016/S2468-2667(20)30088-8)
- Courtney, K. E., & Polich, J. (2009). Binge drinking in young adults: Data, definitions, and determinants. *Psychological Bulletin*, 135(1), 142–156. <https://doi.org/10.1037/a0014414>
- Del Valle-Vera, B., Carmona-Márquez, J., Lozano-Rojas, Ó. M., Parrado-González, A., Vidal-Giné, C., Pautassi, R. M., & Fernández-Calderón, F. (2021). Changes in alcohol use during the COVID-19 pandemic among young adults: The prospective effect of anxiety and depression. *Journal of Clinical Medicine* 2021, 10(19), 4468. <https://doi.org/10.3390/jcm10194468>. Vol. 10, Page 4468.
- El Ansari, W., Salam, A., & Suominen, S. (2020). Prevalence and socio-demographic, academic, health and lifestyle predictors of illicit drug/s use among university undergraduate students in Finland. *International Journal of Environmental Research and Public Health* 2020, 17(14), 5094. <https://doi.org/10.3390/ijerph17145094>. Vol. 17, Page 5094.
- Eurostat. (2021). Alcohol consumption statistics - statistics explained. https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Alcohol_consumption_statistics#General_overview
- Evans, S., Alkan, E., Bhargava, J. K., Tenenbaum, H., & Ng-Knight, T. (2021). Effects of the COVID-19 lockdown on mental health, wellbeing, sleep, and alcohol use in a UK student sample. *Psychiatry Research*, 298, Article 113819. <https://doi.org/10.1016/j.psychres.2021.113819>
- Fernández-Villa, T., Alguacil-Ojeda, J., Ayán-Pérez, C., Bueno-Cavanillas, A., Cancela-Carral, J. M., Capelo-Álvarez, R., Delgado-Rodríguez, M., Jiménez-Mejías, E., Jiménez-Moleón, J. J., Llorca-Díaz, J., Mateos-Campos, R., Molina-De La Torre, A. J., Valero-Juan, L. F., & Martín-Sánchez, V. (2013). Proyecto UNIHOS: cohorte dinámica de estudiantes universitarios para el estudio del consumo de drogas y otras adicciones. *Revista Española de Salud Pública*, 87(6), 575–585. <https://doi.org/10.4321/S1135-57272013000600003>
- Fruehwirth, J. C., Gorman, B. L., & Perreira, K. M. (2021). The effect of social and stress-related factors on alcohol use among college students during the covid-19 pandemic. *Journal of Adolescent Health*, 69(4), 557–565. <https://doi.org/10.1016/j.jadohealth.2021.06.016>
- Gavurova, B., Ivankova, V., & Rigelsky, M. (2021). Alcohol use disorders among Slovak and Czech university students: A closer look at tobacco use, cannabis use and socio-demographic characteristics. *International Journal of Environmental Research and Public Health*, 18(21). <https://doi.org/10.3390/ijerph182111565>
- Gesualdo, C., & Pinquart, M. (2021). Health behaviors of German university freshmen during COVID-19 in association with health behaviors of close social ties, living arrangement, and time spent with peers. *Health Psychology and Behavioral Medicine*, 9(1), 582–599. <https://doi.org/10.1080/21642850.2021.1947291>
- Gogoi, M., Webb, A., Pareek, M., Bayliss, C. D., & Gies, L. (2022). University students' mental health and well-being during the COVID-19 pandemic: Findings from the UniCoVac qualitative study. *International Journal of Environmental Research and Public Health*, 19(15). <https://doi.org/10.3390/ijerph19159322>
- Hennessy, E. A., Tanner-Smith, E. E., Mavridis, D., & Grant, S. P. (2019). Comparative effectiveness of brief alcohol interventions for college students: Results from a network meta-analysis. *Prevention Science*, 20(5), 715–740. <https://doi.org/10.1007/s11211-018-0960-z/TABLES/7>
- Heradstveit, O., Sivertsen, B., Lønning, K. J., & Skogen, J. C. (2022). The extent of alcohol-related problems among college and university students in Norway prior to and during the COVID-19 pandemic. *Frontiers in Public Health*, 10, Article 876841. <https://doi.org/10.3389/fpubh.2022.876841/FULL>
- Holmila, M., & Raitasalo, K. (2005). Gender differences in drinking: Why do they still exist? *Addiction*, 100(12), 1763–1769. <https://doi.org/10.1111/j.1360-0443.2005.01249.X>
- Isaksson, J., Sjöblom, S., Schwab-Stone, M., Stickley, A., & Ruchkin, V. (2019). Risk factors associated with alcohol use in early adolescence among American inner-city youth: A longitudinal study. *Substance Use & Misuse*, 55(3), 358–366. <https://doi.org/10.1080/10826084.2019.1671867>. <https://doi.org/10.1080/10826084.2019.1671867>
- Iwamoto, D. K., & Mui, V. W. (2020). Young adult women and alcohol-related problems: The key role of multidimensional feminine norms. *Substance Abuse: Research and Treatment*, 14. <https://doi.org/10.1177/1178221819888650>
- Jackson, K. M., Merrill, J. E., Stevens, A. K., Hayes, K. L., & White, H. R. (2021). Changes in alcohol use and drinking context due to the COVID-19 pandemic: A multi-method study of college student drinkers. <https://doi.org/10.1111/acer.14574>
- Jaffe, A. E., Kumar, S. A., Ramirez, J. J., & DiLillo, D. (2021). Is the COVID-19 pandemic a high-risk period for college student alcohol use? A comparison of three spring semesters. *Alcoholism: Clinical and Experimental Research*, 45(4), 854–863. <https://doi.org/10.1111/ACER.14572>
- Joinpoint Regression Program, Version 5.1.0.0 – april 2024; statistical methodology and Applications Branch, surveillance research Program, National Cancer Institute.
- Kenney, S. R., DiGuiseppi, G. T., Meisel, M. K., Balestrieri, S. G., & Barnett, N. P. (2018). Poor mental health, peer drinking norms, and alcohol risk in a social network of first-year college students. *Addictive Behaviors*, 84, 151. <https://doi.org/10.1016/j.addbeh.2018.04.012>
- Kilian, C., Allebeck, P., Braddick, F., Gual, A., Barták, M., Bloomeld, K., Neufeld Dresden, M. T., Amy, G. O., Petruzelka, B., Rogalewicz, V., Schulte, B., & Manthey Dresden, J. T. (2021). Alcohol consumption during the COVID-19 pandemic in Europe: A large-scale cross-sectional study in 21 countries (preprint). <https://doi.org/10.21203/rs.3.rs-148341/v2>
- Kilian, C., O'Donnell, A., Potapova, N., López-Pelayo, H., Schulte, B., Miquel, L., Paniello Castillo, B., Schmidt, C. S., Gual, A., Rehm, J., & Manthey, J. (2022). Changes in alcohol use during the COVID-19 pandemic in Europe: A meta-analysis of observational studies. *Drug and Alcohol Review*, 41(4), 918–931. <https://doi.org/10.1111/DAR.13446>
- Killgore, W. D. S., Cloonan, S. A., Taylor, E. C., Lucas, D. A., & Dailey, N. S. (2021). Alcohol dependence during COVID-19 lockdowns. *Psychiatry Research*, 296. <https://doi.org/10.1016/j.psychres.2020.113676>
- Kokotailo, P. K., Egan, J., Gangnon, R., Brown, D., Mundt, M., & Fleming, M. (2004). Validity of the alcohol use disorders identification test in college students. *Alcoholism: Clinical and Experimental Research*, 28(6), 914–920. <https://doi.org/10.1097/01.ALC.0000128239.87611.F5>
- Kosendiak, A., Król, M., Ścisłalska, M., & Kepinska, M. (2022). The changes in stress coping, alcohol use, cigarette smoking and physical activity during covid-19 related lockdown in medical students in Poland. *International Journal of Environmental Research and Public Health*, 19(1). <https://doi.org/10.3390/ijerph19010302/S1>
- Lázaro-Pérez, C., Martínez-López, J. Á., & Gómez-Galán, J. (2020). Addictions in Spanish college students in confinement times: Preventive and social perspective. *Social Sciences* 2020, 9(11), 195. <https://doi.org/10.3390/SOCSCI9110195>. Vol. 9, Page 195.
- López-Moreno, M., Garcés-Rimón, M., Miguel, M., & López, M. T. I. (2021). Adherence to mediterranean diet, alcohol consumption and emotional eating in Spanish university students. *Nutrients* 2021, 13(9), 3174. <https://doi.org/10.3390/NU13093174>. Vol. 13, Page 3174.
- Linden-Carmichael, A. N., Calhoun, B. H., Patrick, M. E., & Maggs, J. L. (2019). University students use fewer protective behavioural strategies on high-intensity drinking days. *Drug and Alcohol Review*, 38(3), 302. <https://doi.org/10.1111/DAR.12913>
- Locke, G. W., Shikret, R., Everett, J. E., & Petry, N. M. (2015). Interpersonal guilt and substance use in college students. *Substance Abuse*, 36(1), 113–118. <https://doi.org/10.1080/08897077.2014.885482>
- Manthey, J., Kilian, C., Carr, S., Bartak, M., Bloomfield, K., Braddick, F., Gual, A., Neufeld, M., O'Donnell, A., Petruzelka, B., Rogalewicz, V., Rossow, I., Schulte, B., & Rehm, J. (2021). Use of alcohol, tobacco, cannabis, and other substances during the first wave of the SARS-CoV-2 pandemic in Europe: A survey on 36,000 European substance users. *Substance Abuse Treatment, Prevention, and Policy*, 16(1), 20. <https://doi.org/10.1186/s13011-021-00373-y>
- Matone, A., Ghirini, S., Gandini, C., & Scafato, E. (2022). Alcohol consumption and COVID-19 in Europe: How the pandemic hit the weak. *Annali Dell'Istituto Superiore Di Sanita*, 58(1), 6–15. https://doi.org/10.4415/ANN.22_01_02
- Mereu, A., Liori, A., Dessi, C., Girau, M., Gilliard, D. C. M., Sotgiu, A., Agabio, R., Contu, P., & Sardù, C. (2021). Alcohol-related behaviour in freshmen university

- students in sardinia, Italy. *International Journal of Environmental Research and Public Health* 2021, 18(13), 7203. <https://doi.org/10.3390/IJERPH18137203>. Vol. 18, Page 7203.
- Messina, M. P., D'angelo, A., Ciccarelli, R., Pisciotto, F., Tramonte, L., Fiore, M., Ferraguti, G., Vitali, M., & Ceccanti, M. (2021). Knowledge and practice towards alcohol consumption in a sample of university students. *International Journal of Environmental Research and Public Health* 2021, 18(18), 9528. <https://doi.org/10.3390/IJERPH18189528>. Vol. 18, Page 9528.
- Millán-Jiménez, A., Herrera-Limones, R., López-Escamilla, Á., López-Rubio, E., & Torres-García, M. (2021). Confinement, comfort and health: Analysis of the real influence of lockdown on university students during the COVID-19 pandemic. *International Journal of Environmental Research and Public Health* 2021, 18(11), 5572. <https://doi.org/10.3390/IJERPH18115572>. Vol. 18, Page 5572.
- Ministerio de Universidades. (2022). Datos y cifras del Sistema Universitario Español. In *Secretaría General Técnica del Ministerio de Universidades*.
- Moure-Rodríguez, L. M., Carbia, C., Varela, M. C., Díaz-Geada, A. D., Cadaveira, F., & Isorna, F. C. (2020). Percepción del consumo de alcohol propio y de allegados en universitarios de primer curso como predictor del consumo a 10 años. *Adicciones*, 35(2), 119–134. <https://doi.org/10.20882/ADICCIONES.1510>
- OEDA. (2021). Observatorio Español de las Drogas y las Adicciones. Encuesta OEDA-COVID: Impacto de la pandemia por Covid-19 durante el año 2020 en el patrón de consumo de sustancias psicoactivas y otros comportamientos con potencial adictivo. Madrid: Ministerio de Sanidad. *Delegación del Gobierno para el Plan Nacional sobre Drogas*; 2021, 42. <https://pnsd.sanidad.gob.es/>.
- OEDA. (2022). Observatorio Español de las Drogas y las Adicciones. EDADES 2022. Encuesta sobre alcohol y otras drogas en España. Madrid: Ministerio de Sanidad. *Delegación del Gobierno para el Plan Nacional sobre Drogas*; 2022, 77.
- Padrón, I., Fraga, I., Vieitez, L., Montes, C., & Romero, E. (2021). A study on the psychological wound of COVID-19 in university students. *Frontiers in Psychology*, 12. <https://doi.org/10.3389/FPSYG.2021.589927>
- Prince, M. A., Read, J. P., & Colder, C. R. (2019). Trajectories of college alcohol involvement and their associations with later alcohol use disorder symptoms. *Prevention Science: The Official Journal of the Society for Prevention Research*, 20(5), 741. <https://doi.org/10.1007/S1121-018-0974-6>
- Puddephatt, J. A., Jones, A., Gage, S. H., Fear, N. T., Field, M., McManus, S., McBride, O., & Goodwin, L. (2021). Associations of alcohol use, mental health and socioeconomic status in England: Findings from a representative population survey. *Drug and Alcohol Dependence*, 219. <https://doi.org/10.1016/J.DRUGALCDEP.2020.108463>
- Real Decreto 463/2020, de 14 de Marzo, Por El Que Se Declara El Estado de Alarma Para La Gestión de La Situación de Crisis Sanitaria Ocasionada Por El COVID-19. BOE 67, de 14 de Marzo, 25390-25400. <https://www.boe.es/boe/dias/2020/03/14/pdfs/BOE-A-2020-3692.pdf>.
- Real Decreto 926/2020, de 25 de Octubre, Por El Que Se Declara El Estado de Alarma Para Contener La Propagación de Infecciones Causadas Por El SARS-CoV-2. BOE 282, de 25 de Octubre, 91072-91084. <https://www.boe.es/boe/dias/2020/10/25/pdfs/BOE-A-2020-12898.pdf>.
- Romero-Blanco, C., Rodríguez-Almagro, J., Onieva-Zafra, M. D., Parra-Fernández, M. L., Prado-Laguna, M. D. C., & Hernández-Martínez, A. (2020). Physical activity and sedentary lifestyle in university students: Changes during confinement due to the COVID-19 pandemic. *International Journal of Environmental Research and Public Health*, 17(18), 1–13. <https://doi.org/10.3390/IJERPH17186567>
- Rosow, I., Bye, E. K., Moan, I. S., Kilian, C., & Bramness, J. G. (2021). Changes in alcohol consumption during the covid-19 pandemic—small change in total consumption, but increase in proportion of heavy drinkers. *International Journal of Environmental Research and Public Health*, 18(8), 4231. <https://doi.org/10.3390/IJERPH18084231/S1>
- Ruiz-Zaldívar, C., García-Garcés, L., Vicario-Merino, Á., Mayoral-Gonzalo, N., Lluésma-Vidal, M., Ruiz-López, M., & Pérez-Manchón, D. (2022). The impact of COVID-19 on the lifestyles of university students: A Spanish online survey. *Healthcare*, 10(2), 309. <https://doi.org/10.3390/HEALTHCARE10020309/S1>
- Saunders, J. B., Aasland, O. G., Babor, T. F., De La Fuente, J. R., & Grant, M. (1993). Development of the alcohol use disorders identification test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction*, 88(6), 791–804. <https://doi.org/10.1111/J.1360-0443.1993.TB02093.X>
- Schmidt, R. A., Genois, R., Jin, J., Vigo, D., Rehm, J., & Rush, B. (2021). The early impact of COVID-19 on the incidence, prevalence, and severity of alcohol use and other drugs: A systematic review. *Drug and Alcohol Dependence*, 228, Article 109065. <https://doi.org/10.1016/J.DRUGALCDEP.2021.109065>
- StataCorp. (2019). StataCorp. 2019. In *Stata statistical software: Release 16*. College Station, TX: StataCorp LLC.
- Steffen, J., Schlichtiger, J., Huber, B. C., & Brunner, S. (2021). Altered alcohol consumption during COVID-19 pandemic lockdown. *Nutrition Journal*, 20(1), 44. <https://doi.org/10.1186/s12937-021-00699-0>
- Tavolacci, M. P., Wouters, E., de Velde, S. Van, Buffel, V., Déchelotte, P., Hal, G. Van, & Ladner, J. (2021). The impact of covid-19 lockdown on health behaviors among students of a French university. *International Journal of Environmental Research and Public Health*, 18(8). <https://doi.org/10.3390/IJERPH18084346>
- Valencia Martín, J. L., Galán, I., Segura García, L., Camarrelles Guillem, F., Suárez Cardona, M., & Brime Beteta, B. (2020). Episodios de consumo intensivo de alcohol "binge drinking": retos en su definición e impacto en salud. *Revista Española de Salud Pública*, 94, 13–14. <https://repisalud.isciii.es/handle/20.500.12105/11474>.
- van Hooijdonk, K. J. M., Rubio, M., Simons, S. S. H., van Noorden, T. H. J., Luijten, M., Geurts, S. A. E., & Vink, J. M. (2022). Student-, study-and COVID-19-related predictors of students' smoking, binge drinking and cannabis use before and during the initial COVID-19 lockdown in The Netherlands. *International Journal of Environmental Research and Public Health*, 19(2), 812. <https://doi.org/10.3390/IJERPH19020812/S1>
- Vasconcelos, M., Crego, A., Rodrigues, R., Almeida-Antunes, N., & López-Caneda, E. (2021). Effects of the COVID-19 mitigation measures on alcohol consumption and binge drinking in college students: A longitudinal survey. *International Journal of Environmental Research and Public Health* 2021, 18(18), 9822. <https://doi.org/10.3390/IJERPH18189822>. Vol. 18, Page 9822.
- White, A., & Hingson, R. (2014). The burden of alcohol use: Excessive alcohol consumption and related consequences among college students. *Alcohol Research: Current Reviews*, 35(2), 201. <https://pubmed.ncbi.nlm.nih.gov/25390712/>.
- White, R., Stevens, K., Hayes, K., & Jackson, K. (2020). Changes in alcohol consumption among college students due to COVID-19: Effects of campus closure and residential change. *Journal of Studies on Alcohol and Drugs*, 81(6), 725–730. <https://doi.org/10.15288/JSD.2020.81.725>. <https://doi.org/10.15288/JSD.2020.81.725>
- WHO, Babor, T. F., Higgins-Biddle, J. C., Saunders, J. B., & Monteiro, M. G. (2001). *Audit: The alcohol use disorders identification test: Guidelines for use in primary health care* (2nd ed.).
- Wicki, M., Kuntsche, E., & Gmel, G. (2010). Drinking at European universities? A review of students' alcohol use. *Addictive Behaviors*, 35(11), 913–924. <https://doi.org/10.1016/J.ADDBEH.2010.06.015>
- Wysokińska, M., & Kołota, A. (2022). Assessment of the prevalence of alcoholic beverage consumption and knowledge of the impact of alcohol on health in a group of polish young adults aged 18–35: A cross-sectional study. *International Journal of Environmental Research and Public Health*, 19(23), Article 15425. <https://doi.org/10.3390/ijerph192315425>
- Yockey, R. A., King, K. A., & Vidourek, R. A. (2020). The epidemiology of recent alcohol use among a national sample of middle eastern college students. *Journal of Drug Education*, 49(1–2), 30–42. <https://doi.org/10.1177/0047237920929328/FORMAT/EPUB>
- Yuan, S., & You, M. (2022). Effects of physical activity on college students' subjective well-being during COVID-19. *Journal of Epidemiology and Global Health*, 12, 441–448. <https://doi.org/10.1007/s44197-022-00062-4>
- Zapata, I., Maté-Muñoz, J. L., Higuera, A., Hernández-Lougedo, J., Martín-Fidalgo, N., García-Fernández, P., Redondo-Vega, M. V., & Ruiz-Tovar, J. (2022). Toxic habits and well-being measures in Spanish healthcare university students during the COVID-19 pandemic. *International Journal of Environmental Research and Public Health* 2022, 19(20), Article 13213. <https://doi.org/10.3390/IJERPH192013213>. Vol. 19, Page 13213.
- Zysset, A., Volken, T., Amendola, S., von Wyl, A., & Dratva, J. (2022). Change in alcohol consumption and binge drinking in university students during the early COVID-19 pandemic. *Frontiers in Public Health*, 10, Article 854350. <https://doi.org/10.3389/FPUH.2022.854350/BIBTEX>